



LIBRARY
OF THE
UNIVERSITY
OF ILLINOIS

630.7

Il 6b

no. 621-640

AGRICULTURE

NON CIRCULATING

**CHECK FOR UNBOUND
CIRCULATING COPY**

330.7
Il 66

LIBRARY OF THE

Bulletin 623

JAN 22 1958

UNIVERSITY OF ILLINOIS
AGRICULTURE LIBRARY.

Performance of EXPERIMENTAL CORN HYBRIDS IN ILLINOIS 1957



LOCATION OF
TEST FIELDS

By R. W. Jugenheimer,
K. E. Williams, and A. J. Crowley, Jr.

CONTENTS

	Page
MATERIAL TESTED	4
MEASURING PERFORMANCE.....	7
RESULTS OF THE TESTS.....	9
NORTHERN ILLINOIS: DeKalb	
Double Crosses (Table 2).....	11
Three-Way, Single, and Double Crosses (Table 3).....	14
Single and Double Crosses (Table 4).....	15
Inbred Lines (Table 5).....	16
NORTH-CENTRAL ILLINOIS: Peoria	
Double Crosses (Table 6).....	17
Single and Double Crosses (Table 7).....	20
Inbred Lines (Table 8).....	21
CENTRAL ILLINOIS: Urbana	
Double Crosses (Tables 9 and 10).....	21, 25
High-Oil Double Crosses and Standards (Table 11).....	26
Three-Way, Single, and Double Crosses (Table 12).....	27
Single and Double Crosses (Table 13).....	28
Corn-Borer-Resistant Single Crosses and Double-Cross Standards (Table 14).....	29
Inbred Lines (Table 15).....	30
Corn-Borer-Resistant Inbred Lines (Table 16).....	30
SOUTH-CENTRAL ILLINOIS: Brownstown	
Double Crosses (Table 17).....	31
Single and Double Crosses (Table 18).....	34
Inbred Lines (Table 19).....	35
SOUTHERN ILLINOIS: Wolf Lake	
Top and Double Crosses (Table 20).....	35
PERFORMANCE OF INBRED LINES IN SINGLE CROSSES (Table 21)...	36
DOUBLE-CROSS HYBRID NUMBERS, PEDIGREES, AND INDEX TO TABLES (Table 22).....	38

Acknowledgment is due W. T. Schwenk and Sons, Edwards, Illinois, and the Shawnee High School, Union county, Illinois, for providing land for two of the tests; to W. C. Jacob and Robert Seif for processing the data; to H. L. Partz, D. R. Browning, and E. Arnzin for conducting the Wolf Lake tests; and to L. H. Lindblom and H. M. Hayes for aid in field and laboratory. Tests in DeKalb, Champaign, and Fayette counties were located on University of Illinois farms managed by R. E. Bell, C. H. Farnham, and P. E. Johnson.

PERFORMANCE OF EXPERIMENTAL CORN HYBRIDS IN ILLINOIS, 1957

By R. W. JUGENHEIMER, K. E. WILLIAMS, and A. J. CROWLEY, JR.¹

ILLINOIS, the center of the corn belt, leads the surrounding states in corn yields per acre. During the ten years 1947-1956, Illinois farmers averaged 55 bushels per acre. These yields have resulted from the use of superior-performing hybrids and modern production practices by efficient seedsmen and farmers. The high yields have brought total production to the point where corn breeders now have the opportunity to reduce the emphasis on yield and to concentrate on developing inbred lines and hybrids that have improved standability, chemical composition, quality, machine harvestability, ear droppage, and resistance to such hazards as insects, diseases, cold, and drouth.

The development and evaluation of superior-performing hybrids is a gradual but continuing procedure. For example, Ill. 960 and Iowa 939 were widely grown in the early days of hybrid corn. These hybrid combinations were completely replaced by such hybrids as Ill. 21 and U.S. 13. These latter hybrids are now rapidly being supplanted by such hybrids as Ill. 1270, Ill. 1570, AES 702, and AES 805. In turn, some of the new experimental Illinois hybrids appear to be superior to these popular combinations.

Illinois hybrids continue to compare favorably with closed-pedigreed hybrids. This is to be expected, of course, since many hybrid seed producers have put out Illinois hybrids under different names or have modified them only slightly by substituting one or two inbred lines. Although many seedsmen use private codes on Illinois hybrids in order to conceal the pedigree of their hybrids, in 1957 some twenty hybrids were grown and certified under their original Illinois designations. Much of this certified seed was grown for sale at wholesale or in interstate commerce; a federal regulation prohibits the assignment of synonyms to corn hybrids used in interstate commerce.

This report summarizes the results of advanced tests of experimental corn hybrids conducted in 1957 by this Station. Data from many preliminary tests involving specialized phases of the corn-research program are not included in this bulletin.

¹R. W. JUGENHEIMER, Professor of Plant Genetics; K. E. WILLIAMS, Fieldman in Agronomy; and A. J. CROWLEY, JR., Research Assistant.

Trials were made at five locations: in DeKalb county in northern Illinois, in Peoria county in north-central Illinois, in Champaign county in central Illinois, in Fayette county in south-central Illinois, and in Union county in southern Illinois. These five locations are representative of the soil, rainfall, and length of growing season in their respective areas.

Hybrids were compared for grain yield, maturity, shelling percentage, standability, ear height and resistance to smut. Only hybrids of similar maturity were tested on the same field. A familiar hybrid whose maturity was considered the standard for the group is named in each table heading.

Since most of the hybrids whose performance is recorded here are not yet in commercial use, the information about them is of most value to producers of hybrid seed. The 1957 performance of hybrids available to farmers in commercial quantities is reported in Bulletin 622 of this Station.

MATERIAL TESTED

Double crosses for consideration of seedsmen. Three hundred and fifty-five different double-cross hybrids were grown at the five locations. Most of the 300 selected Illinois hybrids were developed by the senior author. The seed was produced by controlled hand-pollination.

The double-cross hybrids whose performance is shown in this report and the tables in which each appears are shown in Table 22, which also contains the pedigrees of the hybrids tested. In the pedigrees, the order of the single crosses and of the lines in the single crosses has no significance; it does not indicate which should be used as seed or pollen parent.

Illinois yellow hybrids are numbered consecutively below 2000 and above 3000. White hybrids are numbered in the 2000 series; these white hybrids are usually followed by the letter W. Hybrids that have performed well after regional testing in several corn-belt states have been designated AES (Agricultural Experiment Station) hybrids. Hybrids in the 600 series are similar to Illinois 1277 in maturity; those in the 700 series correspond in maturity to Illinois 21; those in the 800 series correspond to Illinois 1570; and those in the 900 series to Illinois 1851.

The letter A or B following an Illinois hybrid number indicates that the combination of inbred lines making up the hybrid has been rearranged or permuted. For example, if the original pedigree of an Illinois hybrid was (1 × 2) (3 × 4), the letter A following the num-

ber means that the hybrid was put together (1×3) (2×4), the letter B, (1×4) (2×3). A difference in reciprocals is not recognized in this method. When a short dash (-) followed by a number occurs as part of an Illinois hybrid number, it means that a tested related line has been substituted for one of the inbred lines included in the original hybrid.

The University of Illinois does not produce hybrid seed corn in commercial quantities. Hybrids that include new inbred lines may be produced under the "delayed-release" program adopted by the states in the corn belt. Multiplication of a new line is handled by the Station, and the production of single crosses in quantity is handled by the Illinois Seed Producers Association, Champaign, Illinois. If a new Illinois experimental hybrid gives satisfactory performance, the parental lines eventually are released for use by seedsmen.

In order to make the results of corn research more quickly available to the public, the University of Illinois has adopted a slight modification of the "delayed-release" policy as it pertains to Illinois-developed inbred lines. Inbred lines of corn developed by the University of Illinois may be released to the public when they have demonstrated superior combining ability for yield, standability, disease resistance, insect resistance, chemical composition, male sterility, or other characters. Such Illinois lines may form a part of a new hybrid or be used in other ways by corn breeders. Inbred lines of corn developed by others will not be released without their approval.

Hand-pollinated inbred seed of released lines will be available for a fee in packets containing 25 to 100 kernels. Releases will be announced annually on or about April 1. Inquiries may be addressed to the senior author, Agronomy Department, University of Illinois, Urbana, Illinois.

Hybrids for prediction studies. Five sets of single crosses, two sets of three-way crosses, one set of top crosses, and five sets of inbred lines differing in maturity were tested in 1957. The three-way crosses and top crosses (Tables 3, 12, and 20) are a part of the "uniform" tests conducted cooperatively by cornbelt states and the U. S. Department of Agriculture. Seed of the unreleased inbred lines involved in these crosses was contributed by the state or by the federal corn breeder who developed them. Single crosses and inbred lines whose performance is reported in Tables 4, 5, 7, 8, 13, 14, 15, 16, 18, 19, and 21 were developed by the Illinois Station and tested only in Illinois.

The following individuals are responsible at the present time for collecting seed of inbred lines, making the crosses, and distributing

crossed seed of the entries in the cooperative uniform tests: E. C. Rossman (Michigan), D. Linden (Minnesota), N. P. Neal (Wisconsin), and G. H. Stringfield (Ohio) — Table 3; J. H. Lonquist (Nebraska), R. W. Jugenheimer (Illinois), and G. F. Sprague (Iowa) — Table 12; and W. R. Findley (Kansas), F. A. Loeffel (Kentucky), and M. S. Zuber (Missouri) — Table 20.

Performance of single-cross, three-way-cross, and top-cross hybrids is of interest to corn breeders, producers of hybrid seed corn, and farmers. Characteristics of single crosses such as yield, standability, and size, shape, and quality of seed definitely affect the practical production of hybrid seed corn. Some farmers are interested in growing single-cross and three-way-cross hybrids commercially because of their attractive appearance and extreme uniformity. Use of single-cross and three-way-cross data for the prediction of desirable double-cross combinations creates additional interest in the performance of single crosses and three-way crosses.

Prediction studies are an extremely valuable part of a research program. Methods are available to predict the performance of the better hybrid combinations without making and testing large numbers of undesirable crosses. For example, 1,225 single crosses and 690,900 double crosses are possible with 50 inbred lines. However, by using single-cross performance data, the corn breeder can predict which of the many possible double-cross combinations are likely to be most desirable. The following six single crosses can be made with four inbred lines: $A \times B$, $A \times C$, $A \times D$, $B \times C$, $B \times D$, and $C \times D$. The average performance of the four non-parental single crosses gives the predicted performance of a specific double-cross hybrid. For instance, the average yields of the four single crosses $A \times C$, $A \times D$, $B \times C$, and $B \times D$ give the predicted yield of double cross $(A \times B) (C \times D)$. The procedure in predicting acre yields and percentage of erect plants from single-cross data is shown on page 6 of Illinois Agricultural Experiment Station Bulletin 597.

Similar predictions can be made for other characteristics. Predicted hybrid combinations, however, should always be thoroughly tested under field conditions before being put into commercial production.

Three-way crosses also provide useful predictions of the performance of double-cross hybrids. A large number of inbred lines can be compared, and the method is especially valuable where a desirable seed-parent single cross is available for use as a tester. Three-way crosses provide information on specific hybrids and may often eliminate the

time and expense required for testing inbred lines in top crosses and single crosses. The procedure in predicting acre yields and percentage of erect plants from three-way-cross data is also shown on page 6 of Bulletin 597.

Top crosses are simple to produce and often are useful in early stages of a breeding program. For example, a single cross from the corn belt of the United States might contribute genes for high yield and standability, and an open-pollinated variety from Europe might contribute adaptation to local European conditions. Such top crosses might thus combine the desirable traits of the American single cross and the European open-pollinated variety. Most top crosses, however, are temporary expedients, which usually are eventually replaced by double crosses. Top crosses are useful also for evaluating the performance of inbred lines. They also provide a means of selecting promising open-pollinated varieties for use as source material for the development of inbred lines.

MEASURING PERFORMANCE

General information concerning the tests is given in Table 1.

Field plot design. Semi-balanced lattice designs were used to obtain the data reported in Tables 3, 4, 7, 12, 13, and 14. The data in Tables 5, 8, 15, 16, 19, and 20 were obtained in randomized blocks.

Table 1.—GENERAL INFORMATION: Tests of Illinois Experimental Corn Hybrids, 1957

County*	Section of state	Table number	Plants per hill	Date of—	
				Planting	Harvesting
DeKalb.....	Northern	2-5	4	May 7	Oct. 17
Peoria.....	North-Central	6-8	4	May 30	Oct. 29
Champaign.....	Central	9-10	4	June 5	Nov. 7
Champaign.....	Central	11-12	4	June 2	Nov. 4
Champaign.....	Central	13	4	May 8	Oct. 14
Champaign.....	Central	14	4	May 8	Oct. 11
Champaign.....	Central	15-16	4	May 8	Oct. 31
Fayette.....	South-Central	17	3	June 26	Nov. 16
Fayette.....	South-Central	18-19	3	June 26	Nov. 21
Union.....	Southern	20	4	May 2	Oct. 18

* The fields are located near the following cities and towns: in DeKalb county near DeKalb, in Peoria county near Peoria, in Champaign county near Urbana, in Fayette county near Brownstown, and in Union county near Wolf Lake.

Rectangular and simple lattice designs were used for the data reported in Tables 2, 6, 9, 10, 11, 17, and 18. Because of time limitations, the data from the rectangular and simple lattice designs were analyzed by the procedure normally used for randomized block designs.

Method of planting. All plots in these tests were planted, thinned, and harvested by hand in well-fertilized fields prepared in the usual way for corn. Individual plots were 2×5 hills in area. Six kernels were planted in hills spaced 40 inches apart. Hills were thinned to 4 plants at DeKalb, Peoria, Champaign, and Wolf Lake, and to 3 plants at Brownstown.

Acre grain yields. Acre yields are reported as shelled grain containing 15.5 percent moisture, the maximum allowable for No. 2 corn. Data from all plots are included in the report on yield. The only correction for imperfect stands was the following adjustment for missing hills:

$$\text{Ear weight in field} \times \left[1 + \left(\frac{\text{missing hills}}{\text{hills present}} \times .7 \right) \right] = \text{adjusted ear weight}$$

This adjustment adds 0.7 percent of the average hill yield for each missing hill, and assumes that 0.3 percent is made up by the increased yield of surrounding hills.

Shelling percentage and moisture in grain. All ears from one replication of each entry of the double crosses, three-way crosses, and inbred lines were shelled immediately after harvest. Two replications of the single crosses were shelled. The percentage of moisture in the shelled grain was determined with a Steinlite moisture meter.

Stand. Counts of the number of missing hills and number of missing plots were made in late summer in each plot. The data are reported as percentage of a perfect stand. Yields were corrected for missing hills.

Ear height. Representative plants in each plot were measured to determine the distance in inches from the soil to the ear-bearing node.

Erect plants. Percentage of erect plants in each plot of each entry was determined by actual counts at the time of harvest. Stalks broken above the ear were not considered lodged. Stalks leaning less than 45 degrees were considered as erect.

Smutted plants. The number of smutted plants was recorded on all plots in late summer. These data are reported in the tables as percent of smutted plants.

RESULTS OF THE TESTS

Data obtained from the tests are summarized in Tables 2 to 21. Long-time averages are more reliable indexes of the performance of hybrids than a single year's result. The parts of the tables summarizing the results of two or three years therefore deserve the most weight when the results are studied.

Relative performance cannot be determined with absolute accuracy by any method of testing. Small differences between entries are seldom of any significance. In fact, small differences are to be expected among plots planted even with the same lot of seed. Variations in growing conditions such as soil fertility are reduced but not completely eliminated by replicating the same entry several times in the same test. Unavoidable variation may be determined by a mathematical procedure known as analysis of variance. From this procedure figures may be obtained that represent the range which differences between two entries must exceed before those entries can be considered significantly different. The method used to determine this range is called the "Multiple Range Test."¹ This method considers the number of entries that fall within the range as well as the variability of the test. Data shown in **bold-face** were not statistically different from the best performance for that characteristic.

Double-cross hybrids that were high yielding and had excellent standability are indicated by heavy type in Table 22.

The following single crosses, three-way crosses, top crosses, and inbred lines were outstanding in performance in 1957:

Northern Illinois

Table 3A — W136A × (M14 × WF9), W20R × (M14 × WF9), R168 × (M14 × WF9), A257 × (M14 × WF9), MS111 × (M14 × WF9).

Table 3B — MS109 × (WF9 × Oh51A), B47 × (WF9 × Oh51A).

Table 4A — M14 × B14, R113 × B14, R165 × B14, L12 × Oh43, B14 × Oh43, B14 × W64A.

Table 5 — R172, W64A, Oh43, R168.

North-Central Illinois

Table 7A — Hy2 × B14, R109B × B14, R165 × B14, WF9 × B14, B14 × Oh28.

Table 8 — WF9, R168, B14, R172.

¹"Multiple Range and Multiple F Tests," by D. B. Duncan in *Biometrics* 11 (1), 1-43. 1955.

Central Illinois

Table 12A — CI.31A × (Hy × WF9), Oh3F × (Hy × WF9), Oh4G × (Hy × WF9), Oh7N × (Hy × WF9), Oh7P × (Hy × WF9), R168 × (Hy × WF9).

Table 12B — B44 × (WF9 × 38-11), CI.31A × (WF9 × 38-11), Oh3F × (WF9 × 38-11), Oh4G × (WF9 × 38-11), R159 × (WF9 × 38-11), R168 × (WF9 × 38-11).

Table 13A — Hy2 × R71, Hy2 × R74, Hy2 × R127, Hy2 × WF9, R74 × WF9, R129 × WF9, R74 × R129.

Table 14A — R71 × R109B, R74 × R109B, R74 × R112, R74 × R168, R109B × R112, R109B × R114, R112 × R115, R114 × R168, R115 × R168.

Table 15 — R71, R74, 38-11, WF9, R168.

Table 16 — R74, R71, R109B, R113, R114.

South-Central Illinois

Table 18A — C103 × Hy2, C103 × R154, C103 × 38-11, Hy2 × 38-11.

Table 19 — R166, R168, C103.

Southern Illinois

Table 20A — Mo1979 × Mo. 804, Mo9108 × Mo. 804, Ks76-55 × Mo. 804, R159 × Mo. 804, R166 × Mo. 804, CI.90A × Mo. 804, NC220 × Mo. 804.

Results of tests with high-oil hybrids are given in Table 11. Illinois High-Oil hybrids 6052, 6062, 6021, and 6016 were rather outstanding in performance.

Table 2.—DOUBLE CROSSES OF ILLINOIS 1277 MATURITY

Tested in Northern Illinois, 1955-1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Moisture in grain	Shelling	Erect plants	Stand	Ear height	Dropped ears	Smut
A — Three-year averages, 1955-1957									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>
1	Ill. 1555A.....	115	21	79	88	97	45	3	..
2	AES 702.....	114	23	77	90	98	49	7	..
3	Ill. 1863.....	113	23	78	93	98	40	2	..
4	Ill. 1864.....	113	21	78	91	96	41	2	..
5	Ill. 1936.....	113	23	77	92	98	45	2	..
6	Ill. 1281.....	112	22	79	89	99	42	2	..
7	Ill. 1861.....	111	21	79	83	96	43	1	..
8	Ill. 1862.....	111	23	79	91	99	39	2	..
9	ISP 2.....	111	24	77	87	98	44	2	..
10	Ill. 1277.....	110	23	79	86	99	44	2	..
11	Ill. 1375.....	110	22	79	86	97	39	1	..
12	Ill. 1575.....	109	24	77	91	98	45	2	..
13	Ill. 1091A.....	108	23	77	81	93	46	2	..
14	Ill. 1279.....	108	22	79	93	96	43	2	..
15	Ill. 1559B.....	108	23	76	92	97	43	3	..
16	Ill. 1280.....	107	23	78	85	97	43	2	..
17	Ill. 1866.....	107	23	78	92	97	40	1	..
18	Ill. 1289.....	105	23	75	95	98	40	2	..
19	Ohio K24.....	105	21	79	89	94	39	2	..
20	Ill. 1557.....	104	24	77	95	95	42	3	..
21	Ill. 1560A.....	104	23	78	97	97	44	3	..
22	AES 510.....	103	19	79	94	95	41	4	..
23	Ill. 1493.....	101	23	76	94	92	42	2	..
24	Ill. 2247W.....	101	24	77	89	97	46	2	..
25	AES 610.....	99	21	80	92	96	37	1	..
26	Ill. 101.....	97	22	78	85	91	42	3	..
27	Ill. 21.....	90	24	77	82	84	48	2	..
	Average.....	107	22	78	90	96	43	2	..
B — Two-year averages, 1956-1957									
1	AES 702.....	126	22	76	98	97	45	0	0
2	Ill. 1864.....	125	20	79	97	98	42	0	0
3	Ill. 1863.....	122	22	78	97	99	42	0	0
4	Ill. 1956.....	122	22	78	95	98	46	0	0
5	Ill. 1960.....	122	20	79	100	98	45	0	0
6	Ill. 1961.....	122	19	78	99	98	46	0	0
7	Iowa 4757.....	122	20	80	92	98	44	0	1
8	ISP 2.....	122	23	78	87	98	45	0	2
9	Ill. 1281.....	121	22	78	92	100	44	0	0
10	Ill. 1575.....	120	23	78	94	98	45	0	0
11	Ill. 1861.....	120	20	80	90	96	44	0	3
12	Ill. 1862.....	120	23	78	99	100	40	0	2
13	Ill. 1936.....	120	22	78	96	98	46	0	0
14	Ill. 1952.....	120	20	78	97	98	44	0	1
15	Ill. 1957.....	120	20	78	99	98	44	0	2
16	Ill. 1958.....	120	19	80	96	95	46	0	0
17	Ind. 5409.....	120	20	79	94	98	41	0	1
18	Ill. 1277.....	119	23	79	94	99	44	0	1
19	Ill. 1555A.....	119	20	80	96	96	44	0	0
20	Ill. 1955.....	119	20	80	99	96	44	0	0
21	Ill. 1280.....	118	22	78	87	98	46	2	0
22	Ill. 1289.....	118	22	76	98	98	40	0	2
23	Ill. 1559B.....	118	22	76	98	96	43	0	0
24	Ill. 1953.....	118	20	78	98	98	40	0	2
25	Ill. 1962.....	118	20	78	98	98	46	0	1

(Table is continued on next page)

Table 2. — Continued

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Dropped ears	Smut
B — Two-year averages, 1956-1957 — Concluded									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>
26	Ill. 1963.....	118	20	78	98	98	42	0	0
27	Ill. 1091A.....	117	22	78	90	94	46	0	1
28	Iowa 4779.....	117	24	79	98	96	40	0	0
29	Ill. 1375.....	116	22	80	94	96	40	0	0
30	Ill. 1557.....	116	23	76	98	96	43	0	1
31	Ill. 1866.....	116	22	78	94	98	41	0	2
32	Mich. 52-25.....	116	21	80	99	95	38	0	2
33	Minn. CB4621.....	116	18	80	98	98	42	0	1
34	Ill. 1959.....	115	20	79	96	98	42	0	1
35	AES 510.....	114	18	80	98	96	40	0	1
36	Ill. 1279.....	114	21	79	96	96	44	0	2
37	Ill. 1560A.....	114	22	78	100	96	45	1	0
38	Ill. 1493.....	113	22	77	95	91	42	0	2
39	Minn. CB4603.....	112	20	79	100	98	44	0	2
40	Ill. 1902.....	111	23	78	80	91	44	0	2
41	Mich. 53-151.....	111	20	78	96	95	44	0	4
42	Ill. 1954.....	110	20	80	96	94	43	0	2
43	Ohio K24.....	110	20	79	94	96	40	0	2
44	AES 610.....	108	20	80	98	97	38	0	1
45	Ill. 2247W.....	108	23	77	96	95	46	1	0
46	Ill. 101.....	105	21	78	86	88	44	1	2
47	Ill. 21.....	98	23	78	85	78	46	2	2
	Average.....	117	21	78	95	96	43	0	1
C — 1957 results (3 replications)									
1	Ill. 3007.....	137	24	80	99	92	49	0	0
2	Ill. 3009.....	132	22	78	99	98	44	0	3
3	Ill. 3152.....	129	27	78	98	99	42	0	3
4	Ill. 3043.....	128	27	79	100	96	44	0	1
5	Ind. 6225.....	128	22	78	99	99	46	0	1
6	AES 702.....	127	26	75	99	95	45	0	0
7	Ill. 1863.....	127	26	77	99	98	41	0	1
8	Ill. 1952.....	126	24	76	97	98	43	0	1
9	Ill. 3008.....	125	27	78	97	94	46	0	1
10	Ill. 3046.....	125	23	76	98	93	50	0	1
11	Ill. 1864.....	124	23	77	99	96	40	0	1
12	Ill. 1281.....	123	26	77	97	99	43	0	0
13	Ill. 1961.....	123	22	76	99	98	47	0	0
14	Ill. 1999.....	123	31	76	98	98	41	0	0
15	Ill. 1956.....	122	27	75	91	98	47	0	1
16	Ill. 1957.....	122	24	76	100	97	45	0	3
17	Ill. 1862.....	121	27	76	100	100	38	0	2
18	Ill. 1953.....	121	23	76	98	98	41	0	3
19	Ill. 1955.....	121	23	78	100	93	44	0	0
20	Ill. 1960.....	121	25	78	99	98	45	0	0
21	Ill. 3045.....	121	25	79	99	93	42	0	0
22	Ill. 1277.....	120	28	77	93	98	43	0	2
23	Ill. 1575.....	120	29	77	92	96	43	0	0
24	Ill. 1936.....	120	26	75	97	97	44	0	0
25	Ill. 3048.....	120	25	79	100	92	44	0	1
26	Ind. 5409.....	120	25	77	96	96	41	0	1
27	ISP 2.....	120	28	76	84	98	44	0	3
28	Ill. 1091A.....	119	26	77	95	95	45	0	1
29	Ill. 1963.....	119	24	76	98	95	41	0	0
30	Minn. CB4621.....	119	20	77	99	97	40	0	1
31	Ill. 1280.....	118	27	76	87	98	42	0	1
32	Ill. 1555A.....	118	23	77	96	93	45	0	1
33	Ill. 1958.....	118	23	78	98	91	47	0	0
34	Ill. 1959.....	118	25	77	99	98	43	0	2
35	Ill. 1279.....	117	25	77	96	97	45	0	3
36	Ill. 1375.....	117	26	77	94	94	41	0	0
37	Ill. 1962.....	117	24	77	98	97	47	0	2

(Table is concluded on next page)

Table 2.—Concluded

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Dropped ears	Smut
C — 1957 results (3 replications) — Concluded									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>
38	Ill. 1968.....	117	26	76	98	85	50	0	0
39	Ill. 1971.....	117	27	77	95	88	48	0	0
40	Ill. 1998.....	117	26	74	93	95	46	0	1
41	Ill. 3044.....	117	25	75	99	98	45	0	0
42	Iowa 4757.....	117	24	77	95	97	43	0	1
43	Ill. 1559B.....	116	26	74	99	94	43	0	0
44	Ill. 1970.....	116	28	77	91	88	47	0	0
45	Ill. 3047.....	116	25	77	98	88	43	0	0
46	AES 510.....	115	21	77	99	92	41	0	1
47	Ill. 1289.....	115	27	74	97	98	40	0	1
48	Ill. 1861.....	115	24	78	88	91	44	0	4
49	Mich. 52-25.....	115	25	79	99	94	40	0	3
50	Ill. 1560A.....	114	27	77	100	94	45	0	0
51	Iowa 4779.....	114	28	77	97	94	40	0	1
52	Minn. CB4603.....	114	23	77	100	95	43	0	4
53	Ill. 1966.....	113	28	73	94	92	49	0	1
54	Ill. 1866.....	111	26	76	94	97	39	0	2
55	Ill. 3006.....	111	24	77	98	95	41	0	2
56	Ill. 1557.....	110	27	74	99	92	43	0	2
57	Ill. 3016.....	110	30	74	95	93	44	0	2
58	Ill. 3057.....	108	27	77	97	97	45	0	0
59	Ill. 3005.....	107	25	74	97	93	42	0	2
60	Mich. 53-151.....	107	24	77	95	91	43	0	6
61	Ill. 1493.....	106	26	74	94	83	42	0	4
62	Ohio K24.....	106	24	76	96	92	38	0	4
63	AES 610.....	105	24	78	99	96	38	0	1
64	Ill. 1954.....	105	23	78	97	89	43	0	3
65	Mich. 54-70.....	105	26	78	99	88	39	0	1
66	Ill. 1969.....	104	30	77	100	82	46	0	0
67	Mich. 54-116.....	103	24	79	98	99	39	0	5
68	Ill. 1902.....	101	27	76	73	83	44	0	3
69	Ill. 2247W.....	98	28	74	100	92	45	0	1
70	Ill. 101.....	91	25	76	86	79	44	0	4
71	Ohio M15.....	85	24	76	95	76	46	0	1
72	Ill. 21.....	77	28	76	94	58	47	0	2
	Average.....	116	25	77	96	93	44	0	1

Table 3. — THREE-WAY, SINGLE, AND DOUBLE CROSSES
OF ILLINOIS 1277 MATURITY

Tested in Northern Illinois, 1957

(Data in boldface were not statistically different
from the best performance for that characteristic)

Code	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Smut
A — Inbred lines crossed with (M14 × WF9)								
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>
1	Oh26D.....	102	23	76	99	95	34	0
2	W136A.....	107	27	77	97	94	36	1
3	W202.....	107	26	74	98	89	36	8
4	W20R.....	108	25	74	97	90	38	0
5	R165.....	86	31	72	94	81	35	2
6	R168.....	107	26	77	97	84	38	2
7	R172.....	100	29	72	97	90	37	1
8	A257.....	109	26	73	100	98	35	0
9	A296.....	82	22	79	96	82	33	2
10	A568.....	104	24	78	98	89	36	1
11	A569.....	93	25	75	91	89	35	0
12	MS109.....	106	28	75	98	94	38	1
13	MS111.....	119	26	78	98	92	34	2
14	MS121.....	84	29	69	92	79	33	8
15	MS125.....	99	25	73	98	88	36	3
16	MS126.....	104	27	76	99	84	37	1
17	Iowa (Minn.Syn.1).....	105	29	72	73	87	38	1
19	B47.....	110	24	77	85	88	40	3
20	Iowa (Minn.Syn.2).....	97	29	74	97	84	36	0
	Average.....	102	26	75	95	88	36	2
B — Inbred lines crossed with (WF9 × Oh51A)								
21	Oh26D.....	79	24	70	96	89	36	1
22	W136A.....	97	22	79	92	91	35	1
23	W202.....	105	25	77	99	83	38	6
24	W20R.....	102	29	73	96	90	37	4
25	R165.....	95	29	73	95	85	34	0
26	R168.....	100	24	77	99	83	39	2
27	R172.....	88	29	75	98	86	40	2
28	A257.....	99	27	74	99	89	35	1
29	A296.....	105	22	83	93	89	37	4
30	A568.....	100	24	77	97	92	36	1
31	A569.....	103	23	76	97	87	36	1
32	MS109.....	113	28	77	99	97	37	1
33	MS111.....	87	29	76	98	83	35	3
34	MS121.....	91	27	70	96	88	33	1
35	MS125.....	86	27	75	97	85	36	1
36	MS126.....	96	29	77	96	80	38	1
37	Iowa (Minn.Syn.1).....	98	30	69	94	89	39	1
38	Iowa [(M14 × A206) × Oh4C].....	107	26	74	92	92	38	2
39	B47.....	110	24	78	95	91	39	2
40	Iowa (Minn.Syn.2).....	91	31	74	100	90	35	6
	Average.....	98	26	75	96	88	37	2
C — Single crosses								
41	M14 × WF9.....	99	26	73	99	88	33	0
42	WF9 × Oh51A.....	92	26	77	98	76	36	2
	Average.....	96	26	75	98	82	34	1
D — Double crosses								
18	Ill. 3159.....	120	27	76	98	95	39	2
43	Ill. 1277.....	120	28	77	93	93	42	2
44	Ill. 1559B.....	112	30	75	98	93	35	2
45	Ill. 1969.....	107	29	74	97	87	41	2
49	AES 610.....	104	27	73	99	97	37	2
48	Ill. 3059.....	99	29	71	97	92	43	1
47	Ill. 3058.....	96	28	67	100	93	42	0
46	Ill. 3057.....	95	28	67	95	94	39	2
	Average.....	107	28	72	97	93	40	2

Table 4. — SINGLE AND DOUBLE CROSSES
OF ILLINOIS 1277 MATURITY

Tested in Northern Illinois, 1957

(Data in boldface were not statistically different
from the best performance for that characteristic)

Code	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Smut
A — Single crosses								
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>
1	M14×R113.....	108	28	78	98	94	33	0
2	M14×R165.....	91	29	80	56	81	38	0
3	M14×R168.....	114	24	77	93	94	37	0
4	M14×R172.....	114	25	77	100	96	39	0
5	M14×WF9.....	106	27	76	97	90	34	0
6	M14×L12.....	107	25	77	99	98	44	0
7	M14×B14.....	130	24	80	99	91	34	0
8	M14×Oh43.....	112	26	77	100	95	29	0
9	M14×W64A.....	101	26	77	100	95	33	7
12	R113×R165.....	110	29	74	77	91	42	0
13	R113×R168.....	104	25	74	99	93	42	1
14	R113×R172.....	100	28	64	100	99	43	1
15	R113×WF9.....	104	26	73	100	81	39	0
16	R113×L12.....	108	29	74	100	93	48	1
17	R113×B14.....	119	24	76	100	92	45	1
18	R113×Oh43.....	118	32	74	99	96	38	0
19	R113×W64A.....	98	26	71	99	95	38	1
23	R165×R168.....	101	26	75	88	85	43	0
24	R165×R172.....	107	30	75	94	97	44	1
25	R165×WF9.....	121	28	78	88	93	40	0
26	R165×L12.....	88	36	69	30	98	50	3
27	R165×B14.....	119	26	77	92	94	40	0
28	R165×Oh43.....	116	34	78	96	93	36	0
29	R165×W64A.....	103	30	75	74	93	36	1
34	R168×R172.....	73	27	72	98	99	40	0
35	R168×WF9.....	108	24	78	99	84	41	1
36	R168×L12.....	113	29	77	98	97	50	1
37	R168×B14.....	105	24	78	99	90	44	0
38	R168×Oh43.....	104	28	75	98	98	39	0
39	R168×W64A.....	98	23	72	99	95	38	2
45	R172×WF9.....	112	29	76	99	92	40	1
46	R172×L12.....	99	30	70	100	93	45	0
47	R172×B14.....	117	28	77	100	99	45	0
48	R172×Oh43.....	111	34	77	100	93	38	0
49	R172×W64A.....	106	27	79	100	94	38	0
56	WF9×L12.....	89	34	69	97	92	47	1
57	WF9×B14.....	113	25	78	100	83	42	0
58	WF9×Oh43.....	110	32	77	100	92	35	1
59	WF9×W64A.....	65	30	68	91	92	31	2
67	L12×B14.....	108	30	79	99	85	54	0
68	L12×Oh43.....	120	35	78	100	97	45	0
69	L12×W64A.....	112	29	75	94	96	44	3
78	B14×Oh43.....	134	31	77	100	97	37	0
79	B14×W64A.....	123	22	79	100	97	39	1
89	Oh43×W64A.....	117	26	77	100	95	30	1
	Average.....	107	28	75	94	93	40	1
B — Double crosses								
91	Ill. 1277.....	121	28	78	96	95	41	1
93	Ill. 1863.....	113	33	76	99	94	35	1
90	AES 610.....	108	26	74	98	97	34	3
92	Ill. 1555A.....	107	23	79	94	84	42	1
	Average.....	112	28	77	97	92	38	2

Table 5.—INBRED LINES OF ILLINOIS 1277 MATURITY
Tested in Northern Illinois, 1957

(Data in boldface were not statistically different
from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Smut
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>
1	R165.....	48	36	66	55	94	31	1
2	R172.....	47	29	78	100	93	33	0
3	W64A.....	44	23	67	96	86	23	1
4	Oh43.....	41	27	78	100	68	24	0
5	WF9.....	38	34	64	89	84	29	2
6	R168.....	35	21	62	98	93	36	3
7	M14.....	33	25	74	94	75	24	1
8	R113.....	27	25	49	100	77	27	2
9	B14.....	24	34	48	99	91	34	0
10	L12.....	20	40	56	97	66	36	1
	Average.....	36	29	64	93	83	30	1

Table 6. — DOUBLE CROSSES OF ILLINOIS 21 MATURITY

Tested in North-Central Illinois, 1955-1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Moisture in grain	Shelling	Erect plants	Stand	Ear height	Dropped ears	Smut
A — Three-year averages, 1955-1957									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>
1	Ill. 274-1.....	105	20	81	84	96	44	0	..
2	Ill. 972A-1.....	105	21	79	85	97	44	1	..
3	AES 806.....	103	23	79	87	95	39	4	..
4	Ill. 1332.....	103	20	81	86	98	43	2	..
5	Ill. 1912.....	103	21	80	82	94	43	3	..
6	Ill. 1280.....	102	19	81	85	96	40	4	..
7	Ill. 1919.....	101	21	80	84	95	43	2	..
8	AES 805.....	100	22	78	88	95	41	2	..
9	Ill. 1760.....	100	22	79	83	96	39	3	..
10	Ill. 1916.....	100	20	80	82	93	44	3	..
11	Ill. 1511.....	99	21	80	86	96	44	2	..
12	Ill. 1570.....	99	21	78	82	96	44	2	..
13	Ill. 1875.....	99	22	78	89	91	45	6	..
14	Ill. 1575.....	98	21	80	85	97	38	5	..
15	Ill. 1831.....	98	22	80	90	95	37	1	..
16	Ill. 1913.....	98	20	81	82	93	43	5	..
17	Ill. 1917.....	98	21	80	76	91	44	2	..
18	Ill. 1868.....	95	22	79	95	93	39	2	..
19	Ill. 1819.....	94	19	80	84	89	39	1	..
20	Ill. 1863.....	94	20	81	95	94	34	1	..
21	AES 702.....	93	20	79	88	91	41	2	..
22	Ill. 1277.....	93	20	82	87	95	39	1	..
23	Ill. 21.....	92	21	81	86	95	41	1	..
24	Ill. 1555A.....	92	17	80	89	95	38	2	..
25	Iowa 4297.....	92	20	79	87	98	39	1	..
26	Ill. 1560A.....	91	18	82	91	96	38	1	..
27	Ill. 1814.....	90	22	79	94	91	37	1	..
28	Ill. 1873.....	90	20	78	92	94	37	2	..
29	Ill. 2247W.....	86	21	77	86	95	40	3	..
	Average.....	97	21	80	87	94	41	2	..
B — Two-year averages, 1956-1957									
1	AES 805.....	125	20	81	84	96	46	1	2
2	Ill. 972A-1.....	125	18	81	82	97	49	1	2
3	Ill. 1971.....	124	18	84	83	99	46	0	2
4	Ill. 274-1.....	122	18	82	78	95	48	0	1
5	Ill. 1912.....	122	20	81	75	93	48	4	4
6	Ill. 1970.....	122	18	84	79	98	46	0	0
7	AES 806.....	120	21	80	85	93	43	4	1
8	Ill. 1280.....	119	17	82	78	96	45	6	3
9	Ill. 1332.....	119	18	82	84	98	48	2	1
10	Ill. 1511.....	119	18	82	80	96	49	3	0
11	Ill. 1575.....	119	18	82	82	97	42	6	2
12	Ill. 1760.....	119	19	81	76	95	44	3	7
13	Ill. 1570.....	118	19	80	76	96	48	3	3
14	Ill. 1919.....	118	20	81	78	93	48	2	4
15	Ill. 1921.....	118	21	80	93	92	46	1	1
16	Ill. 1928.....	118	22	80	88	94	50	2	4
17	Ill. 1968.....	118	18	82	83	92	48	0	1
18	Ill. 1972.....	118	18	83	82	97	46	0	3
19	Ill. 1973.....	118	19	82	76	97	46	1	2
20	Nebr. 1924.....	117	18	82	90	97	45	2	2
21	Ill. 1831.....	116	20	82	88	94	41	1	3
22	Ill. 1875.....	116	20	80	86	90	50	8	2
23	Ill. 1966.....	116	19	82	86	92	44	0	1
24	Ill. 1969.....	116	18	82	94	94	45	0	2
25	Ill. 1277.....	114	18	84	84	95	42	0	2

(Table is continued on next page)

Table 6. — Continued

Rank in yield	Entry	Acre yield	Moisture in grain	Shelling	Erect plants	Stand	Ear height	Dropped ears	Smut
B—Two-year averages, 1956-1957—Concluded									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>
26	Ill. 1916.....	114	18	81	75	92	50	4	3
27	Iowa 4809.....	114	18	80	95	94	43	1	1
28	Ill. 1913.....	113	18	82	75	92	48	6	0
29	Ill. 1922.....	113	22	80	88	92	46	1	0
30	Ill. 1819.....	112	18	82	78	86	44	0	1
31	Ill. 1868.....	112	20	81	94	94	42	1	1
32	Ill. 1926.....	112	19	80	82	94	46	5	0
33	Ill. 1930.....	112	24	80	88	88	48	2	2
34	AES 702.....	111	18	80	84	88	44	0	0
35	Ill. 21.....	111	18	83	81	93	45	0	2
36	Ill. 1917.....	111	20	81	68	92	50	3	4
37	Iowa 4297.....	110	18	82	84	99	44	0	2
38	Iowa 4879.....	110	18	80	96	90	42	2	1
39	Ill. 1863.....	109	18	82	94	94	38	1	2
40	Ill. 1560A.....	108	17	84	88	96	42	1	2
41	Ill. 1967.....	108	18	80	92	93	48	2	4
42	Ill. 1873.....	107	18	80	90	92	42	2	0
43	Ill. 1927.....	107	20	78	89	90	45	3	0
44	Ill. 1902.....	106	17	82	71	92	44	0	3
45	Ill. 1555A.....	104	15	80	86	92	42	2	2
46	Ill. 1814.....	104	20	82	92	90	40	1	1
47	Ill. 1936.....	104	18	81	91	91	42	1	1
48	Ill. 2247W.....	98	19	78	83	94	44	2	1
49	CB 4726A.....	92	18	82	92	94	38	1	2
	Average.....	114	19	81	84	94	45	2	2
C—1957 results (3 replications)									
1	Ill. 3026.....	127	23	79	85	96	36	0	0
2	Ill. 3027.....	127	22	80	74	98	40	0	0
3	Ill. 3040.....	127	20	79	59	97	46	0	1
4	Ill. 3042.....	127	23	77	92	87	44	0	0
5	Ill. 3033.....	126	19	79	74	98	38	0	3
6	Ill. 3023A.....	125	20	81	87	96	35	0	2
7	Ill. 3035.....	125	22	82	85	94	38	0	0
8	Ill. 3037.....	125	21	80	72	92	43	0	2
9	Ill. 1970.....	123	21	83	63	98	43	0	0
10	Ill. 3010.....	123	22	78	82	98	44	0	2
11	Iowa 4880.....	122	21	79	90	91	39	0	1
12	AES 805.....	121	23	78	79	93	43	0	2
13	Ill. 1332.....	121	21	79	76	99	43	0	1
14	Ill. 1971.....	121	22	82	71	98	43	0	1
15	Ill. 3032.....	121	20	78	89	95	39	0	2
16	Ill. 274-1.....	120	21	80	67	91	47	0	0
17	Ill. 972A-1.....	120	23	78	72	95	47	0	0
18	Ill. 3019.....	120	21	80	88	95	42	0	1
19	Ill. 3014.....	119	24	78	76	94	46	0	1
20	Ill. 3022.....	119	21	81	83	87	40	0	5
21	Ill. 3039.....	118	23	78	88	92	42	0	2
22	Ill. 1280.....	117	20	80	71	97	39	0	1
23	Ill. 1928.....	117	26	76	86	89	50	0	1
24	Ill. 3020.....	117	21	79	87	99	38	0	0
25	Ill. 3029.....	117	21	77	88	88	38	0	0
26	Ill. 1511.....	116	21	79	74	93	45	0	0
27	Ill. 3021.....	116	24	78	93	93	40	0	0
28	Ill. 21.....	115	23	80	79	88	44	0	1
29	Ill. 3012.....	115	23	79	75	91	42	0	3
30	Ill. 3017.....	115	23	79	91	95	39	0	3
31	Ill. 3160.....	115	22	80	92	94	40	0	1
32	Ill. 1575.....	114	20	80	71	94	38	0	1
33	Ill. 1760.....	114	22	79	69	92	39	0	4
34	Ill. 1973.....	114	23	79	64	95	45	0	3
35	Ill. 3043.....	114	20	81	92	93	41	0	0
36	Ill. 3030.....	113	24	78	95	96	39	0	0
37	Ill. 3034.....	113	22	80	82	91	36	0	1

(Table is concluded on next page)

Table 6.—Concluded

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Dropped ears	Smut
C — 1957 results (3 replications) — Concluded									
		<i>bu.</i>	<i>percl.</i>	<i>percl.</i>	<i>percl.</i>	<i>percl.</i>	<i>in.</i>	<i>percl.</i>	<i>percl.</i>
38	Ill. 3169C.....	113	22	74	78	93	36	0	3
39	Ill. 1570.....	112	23	77	67	93	46	0	4
40	Ill. 3013.....	112	25	77	68	95	46	0	0
41	Ill. 3036.....	112	21	80	82	92	34	0	1
42	Nebr. 1924.....	112	21	80	83	94	44	0	2
43	AES 806.....	111	24	78	76	87	38	0	1
44	Ill. 1912.....	111	25	78	67	89	46	0	2
45	Ill. 1966.....	111	22	80	77	87	40	0	1
46	Ill. 1968.....	111	20	80	76	88	45	0	1
47	Ill. 1972.....	111	22	81	74	96	43	0	3
48	Ill. 3018.....	111	22	80	90	94	40	0	3
49	Ill. 3025.....	111	24	80	89	95	40	0	0
50	Ill. 3038.....	111	20	80	91	89	38	0	1
51	Ill. 3041.....	111	19	81	72	93	38	0	3
52	Iowa 4809.....	111	22	78	95	92	41	0	1
53	Ill. 1969.....	110	22	79	93	91	43	0	3
54	Ill. 3045.....	110	20	81	96	89	40	0	0
55	Ill. 3047.....	110	20	78	85	93	41	0	0
56	Ill. 3015A.....	109	22	76	86	100	41	0	0
57	Ill. 3016.....	109	22	79	95	94	37	0	2
58	Ill. 3024.....	109	22	77	96	94	40	0	3
59	CB 4726A.....	109	21	80	87	94	35	0	0
60	Ill. 1831.....	108	23	79	80	89	37	0	2
61	Ill. 1921.....	108	26	77	92	84	45	0	1
62	Ill. 1916.....	107	23	78	71	86	47	0	2
63	Ill. 1926.....	107	24	77	73	93	43	0	0
64	Ill. 1277.....	106	22	81	80	92	40	0	1
65	Ill. 1930.....	106	30	78	86	82	46	0	1
66	Ill. 3011.....	106	24	78	91	98	41	0	0
67	Ill. 3044.....	106	22	78	89	96	40	0	0
68	AES 702.....	104	21	77	71	83	40	0	0
69	Ill. 1560A.....	104	20	83	81	94	39	0	2
70	Ill. 1868.....	104	24	78	93	88	40	0	0
71	Ill. 1875.....	104	24	76	81	81	49	0	0
72	Iowa 4879.....	104	22	78	96	82	38	0	1
73	Ill. 1913.....	103	21	79	65	86	45	0	0
74	Ill. 1967.....	103	23	77	88	89	46	0	4
75	Ill. 1863.....	102	21	80	94	90	32	0	3
76	Ill. 1922.....	102	27	76	84	87	43	0	0
77	Ill. 3124.....	102	21	79	91	88	42	0	1
78	Ill. 1919.....	101	24	78	66	91	46	0	5
79	Ill. 3048.....	101	20	80	92	80	43	0	0
80	Ill. 1902.....	100	20	79	65	87	41	0	3
81	Ill. 1917.....	100	24	77	59	86	49	0	1
82	Iowa 4297.....	98	22	80	75	98	41	0	3
83	Ill. 1927.....	93	24	76	84	81	43	0	0
84	Ill. 1819.....	92	22	78	62	72	41	0	0
85	Ill. 1873.....	91	23	77	82	86	39	0	0
86	Ill. 3046.....	91	20	75	92	83	41	0	0
87	Ill. 1814.....	90	24	78	87	81	36	0	1
88	Ill. 1936.....	90	22	78	89	82	37	0	0
89	Ill. 1555A.....	88	18	73	75	86	39	0	1
90	Ill. 2247W.....	83	23	72	80	89	39	0	0
	Average.....	111	22	79	81	91	41	0	1

Table 7.—SINGLE AND DOUBLE CROSSES
OF ILLINOIS 21 MATURITY

Tested in North-Central Illinois, 1957

(Data in boldface were not statistically different
from the best performance for that characteristic)

Code	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Smut
A—Single crosses								
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>
1	Hy2×R109B.....	109	25	78	67	100	47	1
2	Hy2×R113.....	110	21	79	65	89	47	0
3	Hy2×R165.....	120	21	82	79	90	42	0
4	Hy2×R166.....	93	21	79	66	87	39	0
5	Hy2×R168.....	106	19	81	96	91	42	0
6	Hy2×R172.....	88	21	80	88	77	40	0
7	Hy2×WF9.....	114	22	81	85	90	40	0
8	Hy2×B14.....	118	22	82	87	87	43	1
9	Hy2×Oh28.....	112	19	81	67	86	41	0
12	R109B×R113.....	101	22	78	82	89	37	0
13	R109B×R165.....	108	24	79	78	85	36	1
14	R109B×R166.....	98	24	80	92	94	34	1
15	R109B×R168.....	99	22	82	96	89	39	1
16	R109B×R172.....	90	23	75	81	95	38	1
17	R109B×WF9.....	109	23	77	96	84	34	0
18	R109B×B14.....	130	22	80	92	94	41	0
19	R109B×Oh28.....	91	21	80	87	87	33	1
23	R113×R165.....	106	20	75	46	92	37	0
24	R113×R166.....	79	21	78	47	85	36	1
25	R113×R168.....	94	20	78	86	85	39	0
26	R113×R172.....	92	18	75	86	93	38	0
27	R113×WF9.....	93	19	79	83	78	32	0
28	R113×B14.....	115	19	81	93	93	40	1
29	R113×Oh28.....	79	20	78	89	79	31	0
34	R165×R166.....	98	22	83	42	93	32	0
35	R165×R168.....	107	21	80	87	93	38	1
36	R165×R172.....	86	20	78	90	79	37	0
37	R165×WF9.....	116	20	81	72	93	34	0
38	R165×B14.....	117	21	81	85	90	38	1
39	R165×Oh28.....	97	22	80	65	79	33	1
45	R166×R168.....	82	20	72	78	90	34	2
46	R166×R172.....	82	20	81	63	88	36	1
47	R166×WF9.....	99	20	83	64	89	32	1
48	R166×B14.....	114	19	82	81	100	38	0
49	R166×Oh28.....	82	20	82	42	80	32	3
56	R168×R172.....	59	22	77	88	84	34	0
57	R168×WF9.....	108	21	80	96	94	37	0
58	R168×B14.....	111	18	82	95	88	40	1
59	R168×Oh28.....	94	18	83	88	84	35	1
67	R172×WF9.....	109	21	80	95	88	36	0
68	R172×B14.....	97	18	80	97	84	39	0
69	R172×Oh28.....	86	18	75	77	88	38	3
78	WF9×B14.....	118	20	79	98	88	37	0
79	WF9×Oh28.....	104	19	80	85	87	34	1
89	B14×Oh28.....	120	21	80	92	82	35	1
	Average.....	101	21	79	80	88	37	1
B—Double crosses								
90	AES 702.....	117	22	85	87	84	39	0
92	Ill. 1575.....	108	20	79	85	85	38	1
93	Ill. 1936.....	103	21	75	86	88	34	1
91	Ill. 21.....	97	22	78	70	87	41	1
	Average.....	106	21	79	82	86	38	1

Table 8. — INBRED LINES OF ILLINOIS 21 MATURITY

Tested in North-Central Illinois, 1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Moisture in grain	Shell-ing	Erect plants	Stand	Ear height	Smut
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>
1	WF9	49	19	73	85	87	24	7
2	R168	43	20	76	93	79	30	6
3	Oh28	41	17	69	67	79	29	10
4	B14	40	21	68	100	84	28	0
5	R172	38	24	75	92	84	28	0
6	R165	38	25	58	40	86	29	1
7	R113	35	18	67	95	80	26	0
8	R109B	31	28	69	78	76	31	0
9	Hy2	31	19	68	100	70	30	0
10	R166	17	25	53	78	86	26	41
	Average	36	22	68	83	81	28	6

Table 9. — DOUBLE CROSSES OF ILLINOIS 1570 MATURITY

Tested in Central Illinois (Field A), 1955-1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Moisture in grain	Shell-ing	Erect plants	Stand	Ear height	Dropped ears	Smut	Half-silk
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>	<i>days</i>
A — Three-year averages, 1955-1957										
1	Ill. 1909	117	18	85	65	98	49	6
2	Ill. 1913	116	17	85	64	97	47	3
3	Ill. 1919	116	18	82	66	96	47	1
4	AES 806	114	21	82	70	98	44	5
5	Ill. 1916	114	18	84	70	98	48	4
6	Ill. 274-1	113	18	84	71	93	49	2
7	Ill. 972A-1	113	18	82	78	99	48	5
8	Ill. 1332	113	18	83	76	95	46	6
9	Ill. 1421	113	18	82	69	95	47	2
10	Ill. 1511	112	19	83	72	98	49	8
11	Ill. 1813	112	19	82	79	97	46	8
12	Ill. 1880	112	17	83	72	97	45	4
13	Ill. 1889	112	18	80	77	99	46	3
14	Ill. 1890	112	17	84	79	96	44	7
15	Ill. 1918	112	18	83	68	97	47	4
16	AES 805	111	18	83	80	97	47	5
17	Ill. 1912	109	19	84	59	95	47	4
18	U.S. 13	106	18	82	67	94	52	9
19	Ill. 1570	105	19	82	64	99	49	7
20	AES 808	104	17	84	69	97	43	3
21	Ill. 1767	104	18	80	74	95	45	5
22	Ill. 1777	103	18	82	69	97	45	3
23	Ill. 21	99	17	83	69	96	44	4
	Average	111	18	83	71	97	47	5
B — Two-year averages, 1956-1957										
1	Ind. 5655	131	18	84	92	98	44	1	3	67
2	Ill. 1975	128	21	80	72	97	56	1	3	70
3	Ill. 1913	127	18	85	82	97	48	1	2	65
4	Ill. 1890	126	18	84	89	98	43	3	1	65
5	Ill. 1893	126	18	80	90	93	50	2	4	70
6	Ill. 1909	126	18	84	80	99	48	4	2	65
7	AES 805	124	19	82	90	96	46	2	2	66
8	AES 806	124	22	82	89	98	44	2	4	66
9	Ill. 1332	124	18	82	87	96	46	4	0	66
10	Ill. 1511	124	20	82	88	98	49	5	4	66
11	Ill. 1656	124	18	83	84	100	48	2	2	66
12	Ill. 1916	124	18	84	88	99	48	2	7	66
13	Ill. 1919	124	18	82	81	96	46	0	3	66

(Table is continued on next page)

Table 9. — Continued

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Dropped ears	Smut	Half- silk
B — Two-year averages, 1956-1957 — Concluded										
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>	<i>days</i>
14	Ill. 1928.....	124	20	82	89	96	50	2	4	66
15	Ill. 1889.....	123	18	80	87	100	45	1	6	66
16	Ill. 1421.....	122	19	80	82	96	48	0	1	66
17	Ill. 1813.....	122	20	82	96	96	44	4	1	66
18	Ill. 1918.....	122	18	82	87	98	46	2	1	65
19	Ill. 1921.....	122	19	82	90	98	44	4	0	66
20	Ill. 972A-1.....	121	18	82	92	100	47	2	2	66
21	Ill. 1927.....	121	21	80	90	98	46	2	1	66
22	Ill. 274-1.....	120	18	83	92	94	48	2	1	66
23	Ill. 1880.....	120	17	83	87	98	45	2	1	66
24	Ill. 1973.....	120	19	84	90	100	46	0	2	66
25	Mo. 4060AW.....	120	20	79	93	98	46	2	1	65
26	AES 702.....	119	20	80	86	100	46	3	2	65
27	Ill. 1912.....	119	20	83	71	95	47	4	3	66
28	Iowa 4912.....	119	18	81	84	98	48	2	2	66
29	Ill. 1922.....	118	20	79	94	98	44	3	1	66
30	Ill. 1974.....	118	18	84	84	98	47	0	2	66
31	Ill. 1972.....	117	20	83	81	98	44	0	2	66
32	Ill. 1926.....	116	18	80	92	95	46	4	2	65
33	Ind. 4655.....	116	20	82	96	94	42	2	2	66
34	Ill. 1570.....	115	19	81	72	99	49	2	3	66
35	U.S. 13.....	115	18	82	82	94	50	4	2	66
36	AES 808.....	114	18	84	84	98	42	2	2	67
37	Ill. 1902.....	114	16	83	78	98	44	2	2	66
38	Ill. 1777.....	112	18	82	84	97	44	0	2	66
39	Ill. 1767.....	110	19	80	89	93	45	2	3	66
40	Ind. 4656.....	110	20	82	88	96	41	0	2	66
41	Ill. 21.....	108	18	82	85	96	40	0	0	65
42	Ill. 1935.....	106	18	80	94	96	45	2	6	66
	Average.....	120	19	82	86	97	46	2	2	66
C — 1957 results (3 replications)										
1	Ill. 1332-3.....	119	24	78	84	99	49	0	1	61
2	Ill. 3092.....	118	22	78	90	99	49	0	1	60
3	Ind. 5655.....	118	20	79	87	98	46	0	2	62
4	A 102.....	118	27	78	80	98	45	0	1	62
5	Ill. 1852.....	116	24	76	70	99	50	0	0	66
6	Ill. 1859.....	116	22	78	64	98	51	0	0	65
7	Ill. 1928.....	116	24	80	86	96	52	0	1	61
8	Ill. 1982.....	116	24	76	70	99	52	0	1	62
9	Ill. 1656.....	115	21	80	80	100	47	0	2	62
10	Ill. 1913.....	115	20	82	73	96	49	0	2	60
11	Ill. 1995.....	115	25	88	69	95	47	0	1	62
12	Ill. 3052.....	115	20	80	76	100	44	0	1	59
13	Ill. 3105.....	115	19	81	70	99	49	0	0	60
14	AES 805.....	114	22	78	89	95	47	0	1	60
15	Ill. 1656-2.....	114	21	79	74	94	46	0	2	63
16	Ill. 1890.....	114	20	83	84	97	43	0	0	60
17	Ill. 1975.....	114	25	76	55	96	51	0	1	65
18	Ill. 1981.....	114	22	77	84	98	49	0	2	63
19	Ill. 1985.....	114	21	80	67	99	48	0	1	62
20	Ill. 1991.....	114	24	81	80	92	45	0	2	63
21	Ill. 3049.....	114	21	81	96	94	47	0	0	60
22	Ill. 3091.....	114	25	78	65	99	47	0	1	63
23	Ill. 1921.....	113	22	78	87	98	45	0	0	62
24	Ill. 1976.....	113	24	76	74	91	50	0	1	66
25	Ill. 1978.....	113	25	74	78	98	46	0	0	65
26	Ill. 3093.....	113	22	78	90	99	47	0	0	60
27	Ill. 3095.....	113	21	78	77	96	46	0	0	60
28	Ill. 3104.....	113	20	79	78	96	44	0	1	61
29	AES 702.....	112	23	77	83	99	46	0	0	60
30	Ill. 1656-1.....	112	21	78	83	97	46	0	1	61
31	Ill. 1880.....	112	20	81	81	99	44	0	1	61
32	Ill. 1902.....	112	20	81	69	98	45	0	0	60
33	Ind. 6833.....	112	21	78	80	98	49	0	0	63
34	Ill. 1421.....	111	23	76	71	95	50	0	0	60
35	Ill. 1909.....	111	21	81	73	98	47	0	2	60
36	Ill. 3107.....	111	22	79	82	99	45	0	1	61
37	Ill. 3115.....	111	19	79	78	97	47	0	3	61

(Table is continued on next page)

Table 9.—Continued

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Dropped ears	Smut	Half- silk
C — 1957 results (3 replications) — Continued										
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>	<i>days</i>
38	Ill. 3124	111	24	78	94	99	45	0	2	60
39	AES 806	110	27	78	84	97	44	0	1	61
40	Ill. 21-2	110	21	80	80	90	49	0	1	61
41	Ill. 1332	110	22	78	85	94	44	0	0	61
42	Ill. 1946	110	26	80	85	99	52	0	1	61
43	Ill. 3077	110	20	76	63	98	45	0	3	62
44	Ill. 3083	110	23	80	92	100	44	0	2	60
45	Ill. 3101	110	19	80	68	95	46	0	1	61
46	Mo. 4060AW	110	24	75	90	99	47	0	0	60
47	A 101	110	26	80	80	97	46	0	3	62
48	Ill. 21-4	109	21	79	74	89	43	0	3	60
49	Ill. 1643	109	22	79	71	99	50	0	0	66
50	Ill. 1918	109	22	79	82	100	47	0	1	60
51	Ill. 1974	109	22	81	74	97	49	0	0	61
52	Ill. 1983	109	22	83	88	100	46	0	2	61
53	Ill. 1997	109	25	79	83	97	46	0	0	63
54	Ill. 3062	109	20	82	77	98	48	0	0	62
55	Ill. 3065	109	20	84	76	98	46	0	1	61
56	Ill. 3112	109	22	77	87	92	47	0	2	61
57	Ill. 972A-1	108	22	79	92	99	49	0	0	62
58	Ill. 1332-4	108	23	78	79	99	46	0	2	60
59	Ill. 1337-1	108	22	78	69	97	46	0	0	60
60	Ill. 1851	108	24	75	92	96	49	0	1	66
61	Ill. 1856	108	34	77	73	97	54	0	1	67
62	Ill. 1857	108	29	74	68	98	55	0	2	67
63	Ill. 1927	108	27	77	83	98	46	0	1	61
64	Ill. 1942	108	24	76	81	97	50	0	5	62
65	Ill. 1945	108	27	79	78	99	54	0	1	63
66	Ill. 1973	108	23	82	88	100	47	0	0	61
67	Ill. 1980	108	20	78	90	96	49	0	0	61
68	Ill. 3117	108	22	83	78	98	43	0	3	61
69	Ill. 1332-2	107	21	78	76	96	47	0	0	60
70	Ill. 1570-1	107	23	77	75	100	45	0	1	60
71	Ill. 1813	107	22	78	95	98	45	0	0	60
72	Ill. 1849	107	27	75	88	95	48	0	1	66
73	Ill. 1916	107	23	80	82	98	48	0	8	61
74	Ill. 1984	107	25	77	76	96	46	0	0	61
75	Ill. 3055	107	21	81	84	98	43	0	0	60
76	Ill. 3094	107	25	77	84	97	49	0	4	61
77	Ill. 3114	107	23	77	67	99	49	0	1	60
78	Iowa 4912	107	22	78	76	100	49	0	1	61
79	Ill. 1944	106	23	74	94	97	50	0	3	61
80	Ill. 1988	106	22	78	77	95	44	0	0	61
81	Ill. 1996	106	24	77	86	97	47	0	0	61
82	Ill. 3076	106	22	78	82	98	46	0	0	61
83	Ill. 3116	106	20	80	83	100	45	0	2	61
84	Ill. 3118	106	23	77	85	99	48	0	0	62
85	Ill. 1511	105	24	77	83	98	49	0	2	60
86	Ill. 1570-2	105	24	77	71	99	48	0	0	62
87	Ill. 1889	105	20	76	82	100	45	0	6	61
88	Ill. 1893	105	22	78	85	90	51	0	1	65
89	Ill. 1919	105	22	78	74	100	45	0	2	61
90	Ill. 1922	105	24	75	92	97	45	0	1	61
91	Ill. 3084	105	22	75	82	97	47	0	0	60
92	AES 808	104	21	80	76	99	45	0	2	62
93	Ill. 1570A	104	22	76	73	100	45	0	0	61
94	Ill. 1926	104	20	77	91	94	47	0	1	60
95	Ill. 1986	104	22	78	79	94	44	0	0	60
96	Ill. 1994	104	24	75	95	94	45	0	0	60
97	Ill. 3072	104	23	76	60	98	49	0	0	65
98	Ill. 3089	104	23	78	67	98	48	0	0	61
99	Ill. 3090	104	24	78	68	88	49	0	0	61
100	Ill. 3100	104	20	83	97	97	45	0	3	59
101	Ill. 3121	104	21	80	95	98	44	0	3	60
102	Kan. 1884	104	24	79	66	99	44	0	0	61
103	Ill. 21-3	103	25	79	80	98	48	0	3	61
104	Ill. 1912	103	25	81	55	93	47	0	0	61
105	Ill. 1947	103	23	83	85	95	48	0	1	64
106	Ill. 1972	103	24	81	69	96	45	0	1	61
107	Ill. 1990	103	20	80	84	95	41	0	0	60
108	Ill. 3051	103	22	78	92	97	49	0	1	61
109	Ill. 3056	103	19	80	81	97	43	0	0	60
110	Ill. 3078	103	24	76	70	96	47	0	1	61

(Table is concluded on next page)

Table 9. — Concluded

Rank in yield	Entry	Acre yield	Moisture in grain	Shelling	Erect plants	Stand	Ear height	Dropped ears	Smut	Half-silk
C — 1957 results (3 replications) — Concluded										
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>	<i>days</i>
111	Ill. 3079	103	20	78	76	99	46	0	2	61
112	Ill. 3086	103	22	79	87	95	44	0	3	61
113	Ill. 3098	103	25	79	75	97	47	0	5	60
114	Ill. 3119	103	28	77	86	100	43	0	1	60
115	Ind. 4655	103	24	78	94	90	45	0	1	60
116	Ill. 1925	102	24	80	87	98	43	0	0	60
117	Ill. 1935	102	21	76	88	97	48	0	3	60
118	Ill. 3088	102	20	79	84	92	45	0	0	60
119	Ill. 3102	102	21	78	74	98	48	0	0	60
120	Ill. 3103	102	20	80	74	94	45	0	1	60
121	Ill. 3113	102	19	81	84	98	43	0	0	60
122	Iowa 4906	102	21	81	91	98	44	0	0	61
123	Ill. 21	101	20	80	80	98	41	0	0	60
124	Ill. 1943	101	24	78	87	96	45	0	2	64
125	Ill. 3070	101	25	78	93	98	46	0	1	62
126	Ill. 3080	101	24	78	79	97	48	0	3	61
127	Ill. 3096	101	24	78	90	97	42	0	1	60
128	Ill. 3108	101	20	83	87	90	44	0	3	60
129	Ill. 1777	100	22	78	74	96	46	0	1	61
130	Ill. 1987	100	22	75	82	95	44	0	0	62
131	Ill. 1993	100	20	78	83	98	43	0	0	61
132	Ill. 3067	100	24	85	74	91	45	0	4	63
133	Ill. 3075	100	25	80	92	99	47	0	2	60
134	Ill. 3082	100	23	77	84	92	45	0	3	61
135	Ill. 3087	100	20	81	85	97	43	0	0	60
136	Ill. 3109	100	21	81	79	98	41	0	1	60
137	U.S. 13	100	20	78	76	92	48	0	2	61
138	Ill. 274-1	99	21	81	89	90	48	0	0	61
139	Ill. 1941	99	23	74	76	98	52	0	5	63
140	Ill. 1989	99	21	76	89	92	42	0	0	60
141	Ill. 3085	99	20	79	68	93	48	0	0	61
142	Ill. 1948	98	23	78	78	100	47	0	2	62
143	Ill. 3061	98	20	82	76	96	42	0	0	61
144	Ill. 3064	98	24	80	63	94	45	0	3	61
145	Ill. 3110	98	26	80	82	97	48	0	2	61
146	Ill. 1570	97	23	76	61	99	47	0	2	61
147	Ill. 1977	97	24	78	71	96	48	0	3	61
148	Ind. 4656	97	24	78	81	95	42	0	1	60
149	Ill. 1951	96	22	79	90	93	48	0	1	61
150	Ill. 1992	96	26	73	90	95	47	0	0	61
151	Ill. 3050	96	22	78	82	92	44	0	0	60
152	Ill. 3106	96	25	80	90	98	45	0	2	63
153	Ill. 1767	95	22	75	87	91	45	0	5	61
154	Ill. 1939	95	26	73	92	96	48	0	2	62
155	Ill. 3060	95	25	81	76	95	46	0	1	61
156	Ill. 3081	95	23	79	85	92	44	0	3	61
157	Ill. 3111	95	20	80	89	99	47	0	0	61
158	Ill. 1660	94	37	74	84	100	54	0	0	68
159	Ill. 3058	94	19	80	91	95	46	0	1	61
160	Ill. 3059	94	23	78	84	97	43	0	1	60
161	Ill. 3063	94	23	80	82	93	48	0	2	61
162	Ill. 3069	94	24	78	95	100	44	0	0	62
163	Ill. 3120	94	26	79	71	98	46	0	3	62
164	Ill. 3125	94	26	79	88	97	46	0	2	60
165	Ill. 1949	93	26	77	93	94	47	0	0	63
166	Ill. 3054	93	20	78	78	100	43	0	0	61
167	Ill. 3066	93	23	81	85	92	43	0	1	61
168	Ill. 3097	93	20	78	75	97	44	0	3	61
169	Ill. 1938	92	28	75	95	95	47	0	0	64
170	Ill. 3151	92	21	74	85	95	45	0	0	62
171	Ill. 3071	91	26	75	67	95	50	0	0	65
172	Ill. 3073	91	26	77	94	96	45	0	2	62
173	Ill. 3074	91	23	79	91	98	45	0	1	62
174	Ill. 1950	90	28	77	94	96	45	0	0	64
175	Ill. 3057	90	22	79	85	98	45	0	0	60
176	Ind. 6623	90	20	76	84	98	49	0	1	66
177	Ill. 1940	89	27	75	91	97	46	0	1	63
178	Iowa 4907	89	24	74	90	98	44	0	0	62
179	GCP 6220	88	23	76	90	92	51	0	5	65
180	Ill. 1850	88	34	72	85	96	53	0	1	68
181	Ill. 2246W	73	24	71	78	95	50	0	2	63
	Average	104	23	78	81	97	47	0	1	61

Table 10.—DOUBLE CROSSES OF ILLINOIS 1570 MATURITY
Tested in Central Illinois (Field B), 1956-1957

(Data in boldface were not statistically different
from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Dropped ears	Smut	Half- silk
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>	<i>days</i>
1	Ill. 1981	131	20	80	90	98	49	0	4	68
2	Ill. 1851	130	22	78	93	96	50	0	6	72
3	Ill. 1982	129	22	79	82	98	50	2	2	68
4	Ill. 1995	129	22	86	76	97	49	2	4	67
5	Ill. 1643	128	20	82	82	99	50	3	2	70
6	Ill. 1976	128	22	78	82	95	50	1	4	70
7	Ill. 1978	128	22	78	87	99	47	2	1	69
8	Ill. 1996	128	22	80	91	98	48	1	1	66
9	Ill. 1913	126	18	85	82	97	48	0	3	66
10	Ill. 1991	125	21	80	85	96	46	2	2	69
11	Ill. 1660	124	32	78	88	99	54	0	2	73
12	Ill. 1918	124	20	82	86	99	46	2	1	64
13	Ill. 1945	124	23	82	86	98	54	2	6	68
14	Ill. 1985	124	20	82	80	98	47	2	1	66
15	Ill. 1997	124	22	80	88	98	48	0	1	68
16	Ill. 1332	122	19	80	88	96	46	3	2	66
17	Ill. 1919	122	20	81	84	98	45	0	4	66
18	Ill. 1942	122	22	78	88	98	51	0	10	68
19	Ill. 1944	122	21	78	95	98	50	0	6	66
20	Ill. 1946	122	22	80	88	99	53	0	6	67
21	Ill. 1947	122	22	82	87	97	49	0	3	68
22	Ill. 1980	122	18	81	94	98	48	2	1	66
23	Ill. 1983	122	18	84	92	100	46	2	3	66
24	Ill. 1992	122	22	78	93	97	46	0	1	67
25	Ill. 1984	121	22	78	84	98	47	0	0	66
26	Ill. 1570-1	120	21	80	86	100	46	2	1	66
27	Ill. 1570A	120	20	80	84	100	46	2	0	66
28	Ill. 1880	120	18	83	86	99	45	4	1	66
29	Ill. 1943	120	22	80	92	98	45	0	2	68
30	Ill. 1948	120	22	80	85	100	48	0	5	66
31	Ill. 1994	120	22	78	94	96	46	1	1	66
32	Ill. 1951	119	20	82	91	96	50	0	2	66
33	Ill. 1939	118	24	77	96	98	47	0	4	67
34	Ill. 1941	117	22	78	86	99	51	0	8	68
35	Ill. 1986	117	20	82	89	95	44	0	3	66
36	Ill. 1977	116	20	80	83	98	47	2	4	66
37	Ill. 1987	116	20	78	88	97	46	4	1	67
38	Ill. 1949	115	24	79	96	96	46	0	2	67
39	Ill. 1570	114	20	80	75	98	48	2	2	66
40	Ill. 1988	113	20	80	86	96	44	2	0	66
41	Ill. 1993	113	18	80	90	98	44	1	2	66
42	Ill. 1938	112	26	77	96	97	46	0	2	68
43	Ill. 1990	112	18	82	91	97	40	0	0	64
44	Ill. 1940	110	24	78	94	98	46	0	4	68
45	Ill. 1950	109	25	79	96	98	46	0	0	68
46	Ill. 1989	109	20	79	93	96	42	2	0	66
47	Ill. 2246W	107	20	78	86	98	50	3	2	68
	Average	120	21	80	88	98	47	1	3	67

Table 11.—HIGH-OIL DOUBLE CROSSES AND STANDARDS

Tested in Central Illinois, 1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Oil	Protein	Moisture in grain	Shell-ing	Erect plants	Stand	Ear height	Smut	Half-silk		
A — High-oil double crosses													
		<i>bu.</i>	<i>perct.</i>	<i>lb. per acre</i>	<i>perct.</i>	<i>lb. per acre</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>days</i>
1	Ill. 6062 . . .	137	5.38	413	11.46	879	27	78	55	97	56	1	66
2	Ill. 6052 . . .	136	5.46	416	10.53	802	22	77	67	99	55	1	65
3	Ill. 6016 . . .	131	5.93	435	11.00	807	23	78	50	99	57	3	66
4	Ill. 6021 . . .	131	5.88	431	10.56	774	24	78	60	95	54	1	66
5	Ill. 6084 . . .	124	5.14	357	10.62	738	21	80	25	97	55	1	64
6	Ill. 6076 . . .	122	4.77	326	10.41	711	22	77	23	96	47	1	63
7	Ill. 6075 . . .	119	4.92	328	10.31	687	22	81	45	99	45	1	63
	Average	129	5.35	387	10.70	771	23	78	46	97	53	1	65
B — Standard checks													
1	Ill. 1332 . . .	141	4.43	350	9.46	747	21	80	74	100	48	0	64
2	U.S. 13	137	4.62	354	9.62	738	23	79	65	96	54	1	65
3	K4×38-11 . .	131	5.29	388	11.10	814	28	76	74	97	58	1	69
4	WF9×38-11	126	4.44	313	10.68	754	22	77	79	100	47	2	66
5	Hy2×WF9 . .	122	3.74	255	8.41	575	22	78	78	95	44	1	66
6	Ill. High Oil*	50	11.03	309	12.19	342	20	80	40	90	27	12	30
	Average	118	5.59	328	10.24	662	23	78	68	96	46	3	63

* Open-pollinated variety.

Table 12.—THREE-WAY, SINGLE, AND DOUBLE CROSSES OF ILLINOIS 1570 MATURITY

Tested in Central Illinois, 1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Code	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Smut	Half- silk
A — Inbred lines crossed with (Hy × WF9)									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>days</i>
1	L317.....	137	23	78	36	99	54	0	65
2	38-11.....	134	22	79	47	98	50	0	65
3	B44.....	129	24	81	61	99	43	1	63
4	CI.31A.....	147	27	78	82	99	54	0	65
5	K799.....	134	26	80	83	98	54	1	64
6	K800.....	116	20	80	49	99	50	1	61
7	Mo11662.....	123	26	79	66	97	50	0	64
8	Mo11276.....	122	23	80	62	97	43	1	64
9	Oh3C.....	129	27	76	96	91	46	1	65
10	Oh3F.....	146	25	77	94	99	46	1	64
11	Oh4G.....	135	28	74	91	100	46	0	66
12	Oh7K.....	132	25	78	71	99	46	0	63
13	Oh7N.....	136	24	76	77	100	43	0	63
14	Oh7P.....	144	25	72	82	98	53	1	65
15	R113.....	109	22	77	47	97	44	0	62
16	R153.....	132	24	77	69	100	46	0	62
17	R154.....	127	21	81	51	97	52	2	62
18	R159.....	133	23	80	88	97	48	2	62
19	R166.....	113	22	82	35	99	41	1	62
20	R168.....	130	24	79	98	100	46	2	62
	Average.....	130	24	78	69	98	48	1	63
B — Inbred lines crossed with (WF9 × 38-11)									
21	Hy.....	135	23	78	58	97	53	2	65
22	B44.....	136	21	81	81	97	44	0	65
23	CI.31A.....	139	30	77	82	98	55	1	66
24	K799.....	120	26	77	88	97	47	0	64
25	K800.....	115	21	80	53	97	48	3	64
26	Mo11662.....	129	23	79	73	100	47	2	64
27	Mo11276.....	126	22	78	87	96	44	3	64
28	Oh3F.....	125	28	72	98	98	47	2	66
29	Oh4G.....	140	28	74	86	98	51	1	67
30	Oh7N.....	119	24	75	63	92	45	1	66
31	R113.....	117	26	74	60	93	46	1	65
32	R153.....	119	21	78	63	97	45	1	63
33	R154.....	132	22	80	67	97	51	0	63
34	R159.....	137	22	80	79	99	46	1	64
35	R166.....	110	22	80	48	94	42	4	63
36	R168.....	124	22	79	95	97	46	1	62
	Average.....	126	24	78	74	97	47	1	64
C — Single crosses									
37	Hy×WF9.....	122	22	78	78	95	44	1	60
38	WF9×38-11.....	126	22	77	79	100	47	2	64
	Average.....	124	22	78	78	98	46	2	62
D — Double crosses									
39	Ill. 1332.....	141	21	80	74	100	48	0	64
40	U.S. 13.....	137	23	79	65	96	54	1	65
	Average.....	139	22	80	70	98	51	0	64

Table 13. — SINGLE AND DOUBLE CROSSES
OF ILLINOIS 1570 MATURITY

Tested in Central Illinois, 1957

(Data in boldface were not statistically different
from the best performance for that characteristic)

Code	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Smut	Half- silk
A — Single crosses									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>days</i>
1	Hy2×R71	120	20	81	100	94	45	0	74
2	Hy2×R74	127	19	82	100	98	42	0	72
3	Hy2×R113	114	17	81	98	98	45	1	71
4	Hy2×R127	127	18	86	99	94	44	1	72
5	Hy2×R129	86	19	83	97	91	42	0	75
6	Hy2×R154	118	17	84	91	92	45	0	73
7	Hy2×R168	110	17	82	98	96	43	1	72
8	Hy2×WF9	124	18	84	99	96	40	1	71
9	Hy2×38-11	128	19	84	72	98	49	1	75
12	R71×R74	114	22	81	99	94	37	2	74
13	R71×R113	97	19	80	97	91	41	0	74
14	R71×R127	111	19	83	93	92	40	1	74
15	R71×R129	124	20	80	91	99	42	2	76
16	R71×R154	128	19	84	82	96	43	1	74
17	R71×R168	97	19	83	100	89	41	1	73
18	R71×WF9	120	17	80	97	96	36	0	73
19	R71×38-11	123	18	81	91	96	42	0	75
23	R74×R113	96	18	78	100	92	38	2	71
24	R74×R127	120	19	83	97	99	40	1	73
25	R74×R129	121	20	80	98	93	37	0	72
26	R74×R154	117	18	82	97	92	40	4	71
27	R74×R168	109	18	83	100	99	40	1	71
28	R74×WF9	123	18	81	100	91	33	0	70
29	R74×38-11	118	19	82	99	91	40	1	74
34	R113×R127	107	16	83	95	90	41	1	73
35	R113×R129	109	17	83	100	91	39	1	74
36	R113×R154	107	15	83	93	97	40	1	71
37	R113×R168	93	17	83	99	88	37	3	70
38	R113×WF9	106	16	82	99	94	37	2	70
39	R113×38-11	105	16	82	99	95	40	0	72
45	R127×R129	110	17	83	79	79	40	0	75
46	R127×R154	130	17	85	70	95	43	1	73
47	R127×R168	106	17	85	95	94	40	2	73
48	R127×WF9	110	16	85	96	91	35	1	71
49	R127×38-11	124	18	86	65	86	44	1	75
56	R129×R154	112	16	85	55	89	41	3	73
57	R129×R168	107	19	82	93	89	38	1	73
58	R129×WF9	123	16	82	98	96	36	0	72
59	R129×38-11	112	17	82	86	90	38	2	75
67	R154×R168	111	17	83	93	92	41	2	71
68	R154×WF9	124	17	83	95	94	40	6	71
69	R154×38-11	120	16	86	51	96	47	3	75
78	R168×WF9	107	16	82	98	97	39	0	70
79	R168×38-11	115	16	85	95	97	44	1	73
89	WF9×38-11	111	17	84	96	90	39	1	73
	Average	114	18	83	92	93	41	1	73
B — Double crosses									
91	Ill. 1332	131	18	84	98	94	41	4	74
92	Ill. 1570	128	18	80	94	92	43	8	75
93	Ill. 1893	124	20	81	89	91	42	2	73
90	AES 805	123	18	83	96	96	39	2	71
	Average	126	18	82	94	93	41	4	73

Table 14. — CORN-BORER-RESISTANT SINGLE CROSSES AND DOUBLE-CROSS STANDARDS OF ILLINOIS 1570 MATURITY

Tested in Central Illinois, 1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Code	Entry	Acre yield	Moisture in grain	Shelling	Erect plants	Stand	Ear height	Smut	Half-silk
A — Single crosses									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>days</i>
1	R71×R74	95	20	81	99	96	33	4	73
2	R71×R96B	102	17	85	92	97	33	1	74
3	R71×R109B	107	19	82	98	94	36	4	74
4	R71×R110	94	19	79	98	92	38	3	75
5	R71×R112	93	17	83	98	97	33	5	74
6	R71×R113	96	16	81	100	92	36	1	72
8	R71×R115	100	19	80	93	97	43	11	76
9	R71×R168	109	18	83	95	94	39	1	74
12	R74×R96B	96	16	83	94	93	32	3	72
13	R74×R109B	112	19	81	99	96	36	1	72
14	R74×R110	103	19	78	97	97	34	4	74
15	R74×R112	118	18	84	99	96	33	2	72
16	R74×R113	99	17	81	98	90	38	3	72
17	R74×R114	103	17	79	98	95	36	0	73
18	R74×R115	107	19	80	89	94	42	2	74
19	R74×R168	112	17	84	100	91	37	0	72
23	R96B×R109B	113	17	84	76	94	38	1	74
24	R96B×R110	97	15	82	91	93	36	1	74
25	R96B×R112	104	15	85	98	93	35	4	72
26	R96B×R113	95	15	83	86	95	40	2	73
27	R96B×R114	103	15	86	96	98	35	2	74
28	R96B×R115	95	15	84	81	93	41	5	75
29	R96B×R168	108	16	84	90	89	34	3	74
34	R109B×R110	103	17	80	92	95	38	3	74
35	R109B×R112	121	17	84	98	98	35	0	72
36	R109B×R113	90	16	83	97	91	36	2	73
37	R109B×R114	109	17	81	100	93	38	2	74
38	R109B×R115	118	17	81	83	96	47	2	76
39	R109B×R168	101	17	81	97	90	39	8	72
45	R110×R112	98	15	81	95	94	34	2	73
46	R110×R113	97	15	84	93	94	39	2	75
47	R110×R114	99	15	81	99	95	37	2	75
48	R110×R115	86	15	77	88	98	40	17	75
49	R110×R168	102	16	80	91	92	41	3	73
56	R112×R113	94	16	85	99	91	33	4	72
57	R112×R114	99	15	84	100	94	33	2	74
58	R112×R115	118	18	85	93	90	38	4	74
59	R112×R168	109	17	84	96	95	35	4	73
67	R113×R114	96	15	81	97	92	41	2	74
68	R113×R115	98	16	82	98	94	35	1	72
69	R113×R168	99	15	82	98	92	35	0	71
78	R114×R115	95	14	81	97	93	42	17	75
79	R114×R168	106	16	80	100	90	39	3	73
89	R115×R168	115	16	82	94	94	42	21	75
	Average	103	17	82	95	94	37	4	74
B — Double crosses									
92	AES 805	125	19	82	97	97	36	4	73
91	AES 702	118	18	80	98	92	34	1	72
93	Ill. 3057	106	17	85	100	95	35	0	72
07	Ill. 3059	98	17	79	97	88	37	0	72
90	AES 610	97	16	85	98	93	26	1	69
	Average	109	17	82	98	93	34	1	72

Table 15.—INBRED LINES OF ILLINOIS 1570 MATURITY
Tested in Central Illinois, 1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Mois- ture in grain		Shell- ing	Erect plants	Stand	Ear height	Smut	Half- silk
			<i>bu.</i>	<i>perct.</i>						
1	R71.....	50	18	71	97	97	25	3	82	
2	R74.....	48	18	72	100	94	23	2	80	
3	38-11.....	47	15	77	86	94	33	3	84	
4	R154.....	44	16	77	78	79	31	5	79	
5	WF9.....	41	16	72	99	94	20	7	77	
6	R168.....	39	15	75	82	91	29	4	77	
7	R127.....	38	17	75	79	73	29	1	83	
8	Hy2.....	35	19	70	99	85	27	0	82	
9	R113.....	30	14	60	99	96	24	33	77	
10	R129.....	24	15	67	44	75	23	0	82	
	Average.....	40	16	72	86	88	26	6	80	

Table 16.—CORN-BORER-RESISTANT INBRED LINES
OF ILLINOIS 1570 MATURITY

Tested in Central Illinois, 1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Mois- ture in grain		Shell- ing	Erect plants	Stand	Ear height	Smut	Half- silk
			<i>bu.</i>	<i>perct.</i>						
1	R74.....	50	18	74	100	94	19	1	79	
2	R71.....	47	19	73	97	94	24	3	82	
3	R109B.....	40	17	76	85	86	26	0	80	
4	R168.....	38	16	75	81	90	26	3	77	
5	R113.....	36	14	67	96	89	25	6	76	
6	R114.....	32	15	68	99	90	31	6	86	
7	R112.....	28	16	76	75	92	23	1	80	
8	R110.....	26	14	66	88	87	29	4	84	
9	R96B.....	25	15	69	99	89	22	0	82	
10	R115.....	17	16	62	92	81	34	46	87	
	Average.....	34	16	71	91	89	26	7	81	

Table 17.—DOUBLE CROSSES OF ILLINOIS 1851 MATURITY

Tested in South-Central Illinois, 1955-1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Moisture in grain	Shelling	Erect plants	Stand	Ear height	Dropped ears	Smut
A — Three-year averages, 1955-1957									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>
1	Ill. 1851	81	26	75	66	99	53	0	..
2	AES 805	79	23	77	73	98	49	0	..
3	Ill. 1893	78	25	76	76	100	54	1	..
4	Ill. 1913	78	20	79	71	99	50	1	..
5	AES 904W	77	28	77	69	100	54	0	..
6	Ill. 1349	77	26	79	65	98	55	2	..
7	Ill. 1919	77	22	78	76	96	48	0	..
8	Ill. 1657	76	28	77	46	98	52	0	..
9	Ill. 1918	76	25	77	78	98	48	1	..
10	Ill. 1539A	75	28	77	64	99	54	1	..
11	Ill. 1909	75	22	78	76	96	49	1	..
12	Ill. 1570	74	24	76	68	98	49	1	..
13	Ill. 1332	73	25	77	78	97	50	0	..
14	Ill. 1771	73	27	78	71	95	51	1	..
15	Ill. 1852	73	27	72	63	95	52	0	..
16	AES 903W	72	26	71	59	99	48	0	..
17	Ill. 200	72	26	77	59	98	56	0	..
18	Ill. 1849	72	28	73	76	97	52	2	..
19	Ill. 1850	72	28	76	67	99	53	1	..
20	Ill. 1859	72	27	76	58	99	49	1	..
21	Ill. 1656	71	25	78	66	95	48	0	..
22	Ill. 1856	71	27	73	72	97	51	1	..
23	U.S. 523W	70	27	74	48	98	50	0	..
24	Ill. 1857	69	28	74	55	95	51	1	..
25	Ill. 2246W	69	23	77	66	96	50	1	..
26	Mo. 804	67	25	75	60	100	55	0	..
27	U.S. 13	65	24	76	57	95	51	0	..
	Average	73	26	76	66	98	51	1	..
B — Two-year averages, 1956-1957									
1	Ill. 1851	88	28	74	84	99	56	0	0
2	Ill. 1935	88	24	78	93	99	50	0	0
3	AES 805	86	26	76	90	96	50	0	0
4	Ill. 1919	85	24	77	93	96	48	0	2
5	AES 904W	84	32	76	94	100	54	0	2
6	Ill. 1893	84	28	74	93	100	54	0	0
7	Ill. 1913	84	22	79	95	99	51	1	0
8	Ill. 1349	82	30	78	86	98	57	1	1
9	Ill. 1539A	82	32	76	86	98	55	0	0
10	Ill. 1918	82	28	76	91	96	48	0	0
11	Ill. 1945	82	28	78	88	99	53	0	1
12	Ill. 1948	82	27	77	87	96	50	0	2
13	Ill. 1657	81	30	75	60	98	54	0	0
14	Ill. 1771	80	30	77	90	94	52	1	0
15	Ill. 1889	80	26	76	92	98	49	0	2
16	Ill. 1909	80	26	77	94	96	50	0	0
17	Ill. 1922	80	27	76	96	98	50	0	0
18	Ill. 1928	80	28	75	94	100	50	1	1
19	Ill. 200	79	28	77	80	98	56	0	0
20	Ill. 1332	78	28	76	96	98	52	0	0
21	Ill. 1570	78	26	74	90	98	50	0	0
22	Mo. 916	78	30	72	94	98	54	0	2
23	Ill. 1850	77	30	74	84	100	54	0	1
24	AES 903W	76	30	70	84	98	48	0	0
25	Ill. 1852	76	30	70	80	94	52	0	0
26	Ill. 1859	76	31	74	69	100	48	0	2
27	Mo. 804	76	28	74	81	100	56	0	1
28	Ill. 1849	75	32	68	93	98	53	1	0
29	Ill. 1926	75	26	74	92	97	48	0	0
30	Ill. 1927	75	28	74	90	98	48	0	0

(Table is continued on next page)

Table 17.—Continued

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Dropped ears	Smut
B — Two-year averages, 1956-1957 — Concluded									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>
31	Ill. 1946	75	28	76	90	96	58	0	0
32	Ill. 1656	74	28	76	84	93	48	0	0
33	Ill. 1942	74	30	75	88	100	54	0	2
34	Ill. 1947	74	29	74	90	100	51	0	0
35	Ill. 1951	74	26	78	82	96	55	0	0
36	U.S. 523W	74	30	73	68	98	50	0	1
37	Ill. 1921	73	29	76	84	96	46	0	0
38	Ill. 1944	73	28	74	84	100	52	0	3
39	Ill. 1856	72	30	70	94	96	52	1	2
40	Ill. 1857	72	32	72	69	94	50	0	2
41	Ill. 1940	72	30	74	91	99	52	0	0
42	Ill. 2246W	71	26	76	90	95	50	1	0
43	Ill. 1943	70	28	74	79	96	45	0	0
44	Ill. 1949	70	29	72	96	98	50	0	0
45	Ill. 1939	67	30	72	98	98	48	0	0
46	U.S. 13	67	28	74	75	94	52	0	0
47	Ill. 1938	66	31	70	94	98	48	0	2
48	Ill. 1950	66	28	76	76	96	48	0	0
49	Ill. 1941	64	31	69	92	98	52	0	0
	Average	77	28	75	87	98	51	0	1
C — 1957 results (3 replications)									
1	Ill. 1935	88	35	73	87	99	49	0	0
2	Ill. 1919	87	35	74	93	92	48	0	0
3	Ill. 1851	85	40	67	75	99	55	0	0
4	AES 805	84	37	69	93	96	46	0	0
5	Ill. 2235W	81	39	70	57	99	50	0	0
6	Ill. 1913	80	32	73	92	99	48	0	0
7	Ill. 3136	80	39	73	98	99	47	0	0
8	Ill. 1893	79	40	69	88	99	53	0	0
9	Ill. 1948	79	38	73	75	94	51	0	0
10	Ill. 3126	79	37	72	90	97	50	0	0
11	Ill. 1928	78	41	70	90	99	50	0	0
12	Ill. 1945	78	43	74	80	99	46	0	0
13	Ind. 6874	78	38	66	90	99	50	0	0
14	Ill. 1922	77	38	72	91	97	49	0	0
15	Ill. 3129	77	39	70	78	99	50	0	0
16	Ill. 3133	77	39	73	80	99	48	0	0
17	Ill. 3143	77	39	70	33	99	50	0	0
18	Ill. 3148	77	42	72	98	99	48	0	2
19	Ill. 200	76	41	73	71	97	50	0	0
20	Ill. 1889	76	36	70	90	97	46	0	0
21	Ill. 3147	76	40	73	70	99	51	0	0
22	Ill. 1918	75	40	71	92	94	46	0	0
23	Ill. 1946	75	42	73	81	92	56	0	0
24	Mo. 916	75	44	66	90	96	53	0	0
25	Ill. 1951	74	37	74	67	99	53	0	0
26	Ill. 3149	74	39	73	96	94	48	0	0
27	Ill. 1570	73	38	67	92	97	46	0	0
28	Ill. 1657	73	42	69	33	99	48	0	0
29	Ill. 1771	73	43	74	87	91	50	0	0
30	Ill. 1880	73	37	71	88	97	40	0	0
31	Ill. 1909	73	37	72	100	92	46	0	0
32	Ill. 1947	73	43	71	82	99	47	0	0
33	AES 904W	72	47	69	88	99	48	0	0
34	Ill. 1349	72	42	71	85	97	54	0	0
35	Ill. 1660	72	42	66	59	97	48	0	0
36	Ill. 1927	72	40	71	82	99	42	0	0
37	Ill. 1942	72	42	70	77	99	52	0	0
38	Ill. 1539A	71	48	70	80	97	56	0	0
39	GCP 6220	70	43	71	60	99	54	0	0
40	Ill. 1332	70	41	73	93	96	48	0	0
41	U.S. 619W	70	43	70	70	97	44	0	0
42	Ill. 3138	69	40	71	63	99	56	0	0
43	TRF 13W	69	38	71	75	97	40	0	0

(Table is concluded on next page)

Table 17. — Concluded

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Dropped ears	Smut
C — 1957 results (3 replications) — Concluded									
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>
44	Ill. 1940.....	68	43	69	83	99	50	0	0
45	Ill. 1944.....	68	41	71	72	99	48	0	2
46	Ill. 3059.....	68	36	73	75	99	42	0	0
47	Ill. 3145.....	68	43	69	88	94	46	0	2
48	Ill. 1921.....	67	42	69	72	94	44	0	0
49	U.S. 523W.....	67	45	67	48	99	46	0	0
50	Ill. 1926.....	66	38	69	86	94	44	0	0
51	Ill. 3058.....	66	34	72	97	99	48	0	0
52	Ill. 3128.....	66	44	69	65	99	57	0	0
53	Ill. 3150.....	66	43	67	72	96	46	0	0
54	AES 903W.....	65	43	63	71	97	44	0	0
55	Ill. 1850.....	65	45	65	75	99	48	0	0
56	Ill. 1943.....	65	40	68	61	92	40	0	0
57	Ill. 3139.....	65	42	67	95	99	47	0	0
58	Ill. 3140.....	65	45	65	47	99	56	0	0
59	Ill. 1859.....	64	46	67	52	99	46	0	0
60	Ill. 3130.....	64	43	71	52	94	50	0	0
61	Ill. 3137.....	64	44	68	73	97	55	0	0
62	Mo. 804.....	64	40	68	68	99	52	0	0
63	Ill. 1852.....	63	45	57	69	91	49	0	0
64	Ill. 1950.....	63	41	71	53	99	46	0	0
65	Ill. 3141.....	63	44	71	70	96	55	0	0
66	Mo. 958.....	62	48	61	44	97	54	0	0
67	Ill. 1949.....	61	42	65	91	97	49	0	0
68	Ill. 2246W.....	61	37	68	83	92	48	0	0
69	Mo. 800-3.....	61	46	61	32	99	50	0	0
70	TRF 9W.....	61	45	64	67	99	48	0	3
71	Ill. 1656.....	60	43	69	79	86	40	0	0
72	Ill. 1939.....	60	41	66	97	96	46	0	0
73	Ill. 3057.....	60	38	74	78	92	38	0	0
74	Ill. 3135.....	60	45	65	95	96	47	0	0
75	Ill. 1849.....	59	45	56	90	96	50	0	0
76	Ill. 3144.....	59	47	71	74	91	49	0	0
77	Ill. 3127.....	58	45	66	98	96	48	0	0
78	Ill. 3142.....	58	44	66	54	94	46	0	0
79	Ill. 3146.....	58	42	57	82	99	55	0	0
80	Ind. 6615.....	58	46	61	95	99	46	0	0
81	Ill. 1856.....	57	45	58	97	97	46	0	0
82	Ill. 1938.....	57	45	64	90	96	45	0	0
83	Ill. 1857.....	56	46	65	62	87	46	0	0
84	Ill. 3131.....	56	45	69	79	92	48	0	0
85	TRF 3W.....	56	45	63	73	97	50	0	0
86	U.S. 13.....	55	42	67	61	92	50	0	0
87	TRF 10.....	54	46	68	84	84	42	0	0
88	Ill. 1941.....	53	46	57	84	97	52	0	0
89	Ill. 3132.....	49	47	66	59	84	50	0	0
90	Kan. 2472W.....	48	44	59	71	96	36	0	0
	Average.....	68	42	68	77	96	48	0	0

Table 18. — SINGLE AND DOUBLE CROSSES
OF ILLINOIS 1851 MATURITY

Tested in South-Central Illinois, 1957

(Data in boldface were not statistically different
from the best performance for that characteristic)

Code	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Smut
A — Single crosses								
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>
1	C103×Hy2	94	33	75	95	96	46	0
2	C103×R113	91	30	70	47	97	49	0
3	C103×R153	87	36	71	71	98	49	0
4	C103×R154	97	30	71	88	99	49	0
5	C103×R159	77	36	71	91	97	49	1
6	C103×R166	86	33	76	99	96	40	0
7	C103×R168	88	30	74	84	99	47	0
8	C103×38-11	104	29	72	98	99	53	0
9	C103×Oh7	87	32	69	84	97	46	0
12	Hy2×R113	89	29	71	99	98	48	0
13	Hy2×R153	85	34	75	96	97	44	0
14	Hy2×R154	91	32	78	92	99	47	0
15	Hy2×R159	79	32	73	90	97	49	0
16	Hy2×R166	72	36	77	99	96	38	0
17	Hy2×R168	82	29	77	97	98	44	0
18	Hy2×38-11	93	30	75	93	98	49	0
19	Hy2×Oh7	86	28	78	83	92	47	0
23	R113×R153	73	32	71	88	89	46	0
24	R113×R154	91	29	76	77	99	45	0
25	R113×R159	71	32	71	98	90	47	0
26	R113×R166	72	32	73	78	99	39	0
27	R113×R168	69	29	73	94	98	41	1
28	R113×38-11	87	31	72	81	99	50	0
29	R113×Oh7	74	36	69	46	96	47	0
34	R153×R154	87	31	73	97	99	43	0
35	R153×R159	68	37	67	71	98	49	0
36	R153×R166	68	36	76	76	96	37	0
37	R153×R168	72	34	73	91	99	45	0
38	R153×38-11	76	36	67	59	99	51	0
39	R153×Oh7	70	33	69	93	95	43	0
45	R154×R159	89	30	76	77	97	48	0
46	R154×R166	77	31	80	71	97	42	0
47	R154×R168	87	26	79	98	98	45	0
48	R154×38-11	86	29	76	89	91	52	0
49	R154×Oh7	83	31	77	69	98	44	0
56	R159×R166	70	36	78	99	99	40	0
57	R159×R168	74	30	76	99	95	50	0
58	R159×38-11	90	30	73	84	99	51	0
59	R159×Oh7	70	35	74	82	98	47	0
67	R166×R168	73	29	78	99	99	40	0
68	R166×38-11	71	33	76	95	94	44	0
69	R166×Oh7	71	34	77	99	96	39	0
78	R168×38-11	87	26	78	95	95	49	0
79	R168×Oh7	82	32	79	74	97	44	0
89	38-11×Oh7	78	35	69	64	98	52	0
	Average	81	32	74	86	97	46	0
B — Double crosses								
93	Ill. 1893	97	32	72	74	99	51	1
92	Ill. 1570	93	30	72	78	99	42	0
91	Ill. 1332	92	29	74	93	99	49	1
90	AES 805	83	31	68	89	95	45	0
	Average	91	30	72	84	98	47	0

Table 19. — INBRED LINES OF ILLINOIS 1851 MATURITY
Tested in South-Central Illinois, 1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Rank in yield	Entry	Acre yield	Moisture in grain	Shell-ing	Erect plants	Stand	Ear height	Smut
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>
1	R166.....	37	23	75	75	95	20	1
2	Oh7.....	32	33	72	40	97	31	0
3	R159.....	26	42	65	35	82	33	0
4	R153.....	25	42	63	48	82	30	0
5	R168.....	24	42	71	72	92	29	0
6	C103.....	21	34	63	71	74	28	0
7	38-11.....	19	42	62	55	98	34	0
8	R154.....	19	42	60	40	91	33	0
9	Hy2.....	16	42	54	44	88	27	0
10	R113.....	12	42	45	34	93	26	0
	Average.....	23	38	63	51	89	29	0

Table 20. — TOP AND DOUBLE CROSSES
OF ILLINOIS 1851 MATURITY

Tested in Southern Illinois, 1957

(Data in boldface were not statistically different from the best performance for that characteristic)

Code	Entry	Acre yield	Moisture in grain	Shell-ing	Erect plants	Stand	Ear height
A — Inbred lines crossed with (Mo. 804)							
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>
1	Mo1979.....	100	16	80	99	95	50
2	Mo9108.....	100	17	85	99	98	48
3	Mo9294.....	92	16	84	100	97	53
4	Mo11077.....	89	16	89	99	99	42
5	Mo11144.....	105	16	83	97	98	47
6	Mo11153.....	95	16	85	99	96	48
7	Mo11276.....	75	16	80	99	99	39
8	Mo01392.....	86	17	86	100	97	45
9	Mo01480.....	92	16	85	96	97	47
10	Ks49-55.....	88	16	83	98	96	55
11	Ks54-55.....	83	17	75	99	94	44
12	Ks76-55.....	113	17	91	100	99	40
13	Ks86-55.....	81	16	86	94	98	49
14	Ks133-53.....	100	16	85	96	98	46
15	Ks159-53.....	78	16	85	100	100	48
16	Oh8.....	85	17	85	100	96	47
17	R113.....	78	15	83	99	98	46
18	R153.....	92	16	83	99	96	49
19	R154.....	93	15	84	87	84	42
20	R159.....	104	15	85	98	96	45
21	R166.....	98	16	89	99	97	38
22	R173.....	84	15	83	99	99	36
23	C1.90A.....	100	17	85	99	98	47
24	C1.91B.....	76	16	84	98	96	55
25	NC218.....	93	18	84	100	96	50
26	NC220.....	111	17	84	99	94	51
27	NC222.....	86	20	81	100	97	54
28	NC224.....	93	16	83	100	98	42
	Average.....	92	16	84	98	97	47
B — Double crosses							
30	Ill. 1851.....	119	15	85	99	98	50
29	Mo. 804.....	108	15	84	99	99	48
	Average.....	114	15	84	99	98	49

Table 21.—AVERAGE PERFORMANCE OF INBRED LINES
AS MEASURED IN SINGLE CROSSES^a

(Comparisons can be made only within each section)

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Dropped ears	Smut	Half- silk
A — Ill. 1277 maturity (summarized from Table 4)										
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>	<i>days</i>
1	B14.....	119	26	78	99	92	42	0	0	..
2	Oh43.....	116	31	77	99	95	36	0	0	..
3	M14.....	109	26	78	94	93	36	0	1	..
4	R113.....	108	27	73	97	93	41	0	1	..
5	R165.....	106	30	76	77	92	41	0	1	..
6	L12.....	105	31	74	91	94	47	0	1	..
7	R172.....	104	29	74	99	96	41	0	0	..
8	W64A.....	103	27	75	95	95	36	0	2	..
9	WF9.....	103	28	75	97	89	39	0	1	..
10	R168.....	102	26	75	97	93	42	0	1	..
	Average.....	108	28	76	94	93	40	0	1	..
B — Ill. 21 maturity (summarized from Table 7)										
1	B14.....	116	20	81	91	90	39	0	1	..
2	Hy2.....	108	21	80	78	89	42	0	0	..
3	WF9.....	108	21	80	86	88	35	0	0	..
4	R165.....	106	21	80	72	88	36	0	0	..
5	R109B.....	104	23	79	86	91	38	0	1	..
6	R113.....	97	20	78	75	87	37	0	0	..
7	R168.....	96	20	79	90	89	38	0	1	..
8	Oh28.....	96	20	80	77	84	35	0	1	..
9	R166.....	92	21	80	64	90	35	0	1	..
10	R172.....	88	20	78	85	86	37	0	1	..
	Average.....	101	21	80	80	88	37	0	1	..
C — Ill. 1570 maturity (summarized from Table 13)										
1	R154.....	119	17	84	81	94	42	0	2	72
2	Hy2.....	117	18	83	95	95	44	0	1	73
3	38-11.....	117	17	84	84	93	43	0	1	74
4	R74.....	116	19	81	99	94	39	0	1	72
5	R127.....	116	17	84	88	91	41	0	1	73
6	WF9.....	116	17	83	98	94	37	0	1	71
7	R71.....	115	19	81	94	94	41	0	1	74
8	R129.....	112	18	82	89	91	39	0	1	74
9	R168.....	106	17	83	97	93	40	0	1	72
10	R113.....	104	17	82	98	93	40	0	1	72
	Average.....	114	18	83	92	93	41	0	1	73

^a Calculated for each inbred by averaging the performance of single crosses in which it was one of the parents.

(Table is concluded on next page)

Table 21.—Concluded

Rank in yield	Entry	Acre yield	Mois- ture in grain	Shell- ing	Erect plants	Stand	Ear height	Dropped ears	Smut	Half- silk
D — Ill. 1570 maturity (summarized from Table 14)										
		<i>bu.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>perct.</i>	<i>in.</i>	<i>perct.</i>	<i>perct.</i>	<i>days</i>
1	R109B.....	108	17	82	93	94	38	0	3	73
2	R168.....	107	16	82	96	92	38	0	5	73
3	R112.....	106	16	84	97	94	34	0	3	73
4	R74.....	105	18	81	97	94	36	0	2	73
5	R115.....	104	17	81	91	94	41	0	9	75
6	R96B.....	101	16	84	89	94	36	0	2	74
7	R114.....	101	16	82	98	94	38	0	4	74
8	R71.....	100	18	82	97	95	36	0	4	74
9	R110.....	98	16	80	94	94	37	0	4	74
10	R113.....	96	16	82	96	92	37	0	2	73
	Average.....	103	17	82	95	94	37	0	4	74
E — Ill. 1851 maturity (summarized from Table 18)										
1	C103.....	90	32	72	84	98	48	0	0	..
2	R154.....	88	30	76	84	97	46	0	0	..
3	Hy2.....	86	31	75	94	97	46	0	0	..
4	38-11.....	86	31	73	84	97	50	0	0	..
5	R113.....	80	31	72	79	96	46	0	0	..
6	R168.....	79	29	76	92	98	45	0	0	..
7	Oh7.....	78	33	73	77	96	45	0	0	..
8	R153.....	76	34	71	82	97	45	0	0	..
9	R159.....	76	33	73	88	97	48	0	0	..
10	R166.....	73	33	77	91	97	40	0	0	..
	Average.....	81	32	74	86	97	46	0	0	..

Table 22. — DOUBLE-CROSS HYBRID NUMBERS,
PEDIGREES, AND INDEX TO TABLES

(Hybrids that were high yielding and had excellent standability are indicated
by table numbers in boldface type)

Hybrid	Pedigree	Table No.
Illinois hybrids		
21	(Hy2 × 187-2) (WF9 × 38-11)	2ABC, 6ABC, 7B, 9ABC
21-2	(HyR × 187R) (WF9TMS × 38-11)	9C
21-3	(WF9 × 38-11) (187-2 × Cl.42A)	9C
21-4	(HyR × 187-2) (WF9TMS × 38-11)	9C
101	(M14 × WF9) (187-2 × W26)	2ABC
200	(WF9 × 38-11) (L317 × K4)	17ABC
274-1	(Hy2 × WF9) (Oh7 × 187-2)	6ABC, 9ABC
972A-1	(Hy2 × L317) (WF9 × Oh7)	6ABC, 9ABC
1091A	(Hy2 × 187-2) (M14 × WF9)	2ABC
1277	(M14 × WF9) (L205 × 187-2)	2ABC, 3D, 4B, 6ABC
1279	(M14 × WF9) (A375 × 187-2)	2ABC
1280	(M14 × WF9) (Os420 × 187-2)	2ABC, 6ABC
1281	(M14 × WF9) (A374 × A375)	2ABC
1289	(M14 × W22) (WF9 × L205)	2ABC
1332	(Hy2 × Oh7) (WF9 × 38-11)	6ABC, 9ABC, 10, 11B, 12D, 13B, 17ABC, 18B
1332-2	(HyR × Oh7R) (WF9TMS × 38-11)	9C
1332-3	(WF9 × 38-11) (Oh7 × Cl.42A)	9C
1332-4	(HyR × Oh7) (WF9TMS × 38-11)	9C
1337-1	(HyR × R61) (WF9TMS × 38-11)	9C
1349	(38-11 × Mo940) (K155 × K201)	17ABC
1375	(M14 × WF9) (N6 × Oh51A)	2ABC
1421	(Hy2 × WF9) (P8 × Oh7)	9ABC
1493	(WF9 × L205) (Oh28 × W22)	2ABC
1511	(Hy2 × WF9) (38-11 × L304A)	6ABC, 9ABC
1539A	(38-11 × Cl.7) (K201 × Cl.21E)	17ABC
1555A	(WF9 × Oh51A) (L224 × Oh28)	2ABC, 4B, 6ABC
1557	(M14 × Oh28) (L205 × Oh51A)	2ABC
1559B	(M14 × Oh28) (WF9 × Oh51A)	2ABC, 3D
1560A	(WF9 × Oh51A) (L205 × Oh28)	2ABC, 6ABC
1570	(Hy2 × Oh41) (WF9 × 38-11)	6ABC, 9ABC, 10, 13B, 17ABC, 18B
1570A	(Hy2 × WF9) (38-11 × Oh41)	9C, 10
1570-1	(HyR × Oh41) (WF9TMS × 38-11)	9C, 10
1570-2	(WF9 × 38-11) (Oh41 × Cl.42A)	9C
1575	(M14 × WF9) (L12 × Oh28)	2ABC, 6ABC, 7B
1643	(Cl03 × 38-11) (Hy2 × Oh7)	9C, 10
1656	(Cl03 × Hy2) (WF9 × 38-11)	9BC, 17ABC
1656-1	(Cl03 × HyR) (WF9TMS × 38-11)	9C
1656-2	(Cl03 × Cl.42A) (WF9 × 38-11)	9C
1657	(K4 × Oh7) (K201 × Cl.21E)	17ABC
1660	(K4 × K201) (Oh7 × Cl.21E)	9C, 10, 17C
1760	(WF9 × 38-11) (Oh29 × Oh45)	6ABC
1767	(Hy2 × Oh45) (WF9 × 38-11)	9ABC
1771	(Oh7B × Cl.7) (T8 × Cl.21E)	17ABC
1777	(Hy2 × WF9) (R114 × R116)	9ABC
1813	(Cl03 × Oh45) (Hy2 × WF9)	9ABC
1814	(Hy2 × WF9) (M14 × Oh45)	6ABC
1819	(R2 × WF9) (R61 × Oh43)	6ABC
1831	(WF9 × W146) (K237 × Oh45)	6ABC
1849	(Cl03 × 38-11) (K201 × Cl.21E)	9C, 17ABC
1850	(Cl03 × Cl.21E) (38-11 × K201)	9C, 17ABC

(Table is continued on next page)

Table 22. — Continued

Hybrid	Pedigree	Table No.
Illinois hybrids (continued)		
1851	(C103 × 38-11) (Oh7 × Cl.21E)	9C, 10, 17ABC, 18B
1852	(C103 × Cl.21E) (38-11 × Oh7)	9C, 17ABC
1856	(38-11 × Oh7) (K201 × Cl.21E)	9C, 17ABC
1857	(38-11 × Oh41) (K201 × Cl.21E)	9C, 17BC
1859	(38-11 × Oh7) (Oh41 × Cl.21E)	9C, 17ABC
1861	(M14 × WF9) (I.224 × Oh28)	2ABC
1862 (Iowa 4779)	(M14 × WF9) (Oh43 × Oh51A)	2ABC
1863	(M14 × WF9) (I.205 × Oh43)	2ABC, 4B, 6ABC
1864	(M14 × WF9) (Oh43 × W22)	2ABC
1866	(M14 × WF9) (Oh26A × Oh45)	2ABC
1868	(C103 × Oh43) (Hy2 × WF9)	6ABC
1873	(C103 × M14) (R75 × Oh43)	6ABC
1875	(C103 × 38-11) (Hy2 × WF9)	6ABC
1880	(R103 × R104) (WF9 × 38-11)	9ABC, 10, 17C
1889	(C103 × Oh45) (38-11 × Oh29)	9ABC, 17BC
1890	(C103 × Oh45) (R75 × 38-11)	9ABC
1893	(C103 × 38-11) (Oh7B × Oh29)	9BC, 13B, 17ABC, 18B
1902	(R138 × R142) (R139 × R141)	2BC, 6BC, 9BC
1909	(R130 × R151) (WF9 × 38-11)	9ABC, 17ABC
1912	(R151 × R156) (WF9 × 38-11)	6ABC, 9ABC
1913	(R151 × R154) (WF9 × 38-11)	6ABC, 9ABC, 10, 17ABC
1916	(R130 × R154) (WF9 × 38-11)	6ABC, 9ABC
1917	(R153 × R154) (WF9 × 38-11)	6ABC
1918	(R151 × R153) (WF9 × 38-11)	9ABC, 10, 17ABC
1919	(R130 × R156) (WF9 × 38-11)	6ABC, 9ABC, 10, 17ABC
1921	(R71 × R105) (WF9 × 38-11)	6BC, 9BC, 17BC
1922	(Hy2 × WF9) (R71 × R105)	6BC, 9BC, 17BC
1925	(Hy2 × WF9) (R71 × R113)	9C
1926	(R71A × R74) (R75 × 38-11)	6BC, 9BC, 17BC
1927	(Hy2 × WF9) (R71A × R74)	6BC, 9BC, 17BC
1928	(R75 × 38-11) (R98 × R105)	6BC, 9BC, 17BC
1930	(Hy2 × WF9) (R98 × R105)	6BC
1935	(C103 × R101) (R75 × 38-11)	9BC, 17BC
1936	(Hy2 × WF9) (M14 × B14)	2ABC, 6BC, 7B
1938	(R71 × R105) (R98 × R153)	9C, 10, 17BC
1939	(R71 × R98) (R105 × R153)	9C, 10, 17BC
1940	(R71 × R153) (R98 × R105)	9C, 10, 17BC
1941	(R98 × R105) (R130 × R153)	9C, 10, 17BC
1942	(R98 × R153) (R105 × R130)	9C, 10, 17BC
1943	(R71 × R105) (R153 × R154)	9C, 10, 17BC
1944	(R71 × R98) (R130 × R153)	9C, 10, 17BC
1945	(R98 × R151) (R105 × R130)	9C, 10, 17BC
1946	(R98 × R155) (R105 × R130)	9C, 10, 17BC
1947	(R105 × R130) (R153 × R155)	9C, 10, 17BC
1948	(R105 × R151) (R153 × R154)	9C, 10, 17BC
1949	(R71 × R105) (R151 × R153)	9C, 10, 17BC
1950	(R71 × R105) (R153 × R155)	9C, 10, 17BC
1951	(R71 × R130) (R98 × R155)	9C, 10, 17BC
1952	(M14 × B14) (A545 × W64A)	2BC
1953	(M14 × A223) (B14 × W64A)	2BC
1954	(M14 × A297) (B14 × A545)	2BC
1955	(M14 × A297) (B14 × W64A)	2BC
1956	(M14 × A545) (B14 × A239)	2BC
1957	(M14 × A545) (B14 × W64A)	2BC
1958	(M14 × Oh26A) (B14 × A545)	2BC

(Table is continued on next page)

Table 22. — Continued

Hybrid	Pedigree	Table No.
Illinois hybrids (continued)		
1959 (Ind. 6225)	(M14 × W64A) (B14 × A297)	2BC
1960	(M14 × W64A) (B14 × A545)	2BC
1961	(B14 × A545) (A239 × W64A)	2BC
1962	(B14 × A545) (A297 × W64A)	2BC
1963	(B14 × A545) (Oh26A × W64A)	2BC
1966	(R163 × R165) (WF9 × B14)	2C, 6BC
1967	(R163 × R168) (WF9 × B14)	6BC
1968	(R163 × R169) (WF9 × B14)	2C, 6BC
1969	(R165 × R168) (WF9 × B14)	2C, 3D, 6BC
1970	(R165 × R169) (WF9 × B14)	2C, 6BC
1971	(R168 × R169) (WF9 × B14)	2C, 6BC
1972	(R163 × R165) (R168 × R169)	6BC, 9BC
1973	(R163 × R168) (R165 × R169)	6BC, 9BC
1974	(R163 × R169) (R165 × R168)	9BC
1975	(WF9 × Cl.38B) (Cl.42A × Cl.317B)	9BC
1976	(38-11 × Oh41) (Oh7 × Cl.21E)	9C, 10
1977	(WF9 × 38-11) (Oh29 × Oh41)	9C, 10
1978	(Cl03 × 38-11) (WF9 × Oh7A)	9C, 10
1980	(Cl03 × B14) (WF9 × 38-11)	9C, 10
1981	(WF9 × 38-11) (Oh7 × Cl.21E)	9C, 10
1982	(Cl03 × 38-11) (WF9 × Cl.21E)	9C, 10
1983	(Hy2 × B14) (WF9 × 38-11)	9C, 10
1984	(Hy2 × WF9) (Oh29 × Oh41)	9C, 10
1985	(Hy2 × WF9) (R61 × Oh41)	9C, 10
1986	(Hy2 × WF9) (Oh43 × 187-2)	9C, 10
1987	(Cl03 × B10) (Hy2 × WF9)	9C, 10
1988	(Cl03 × R61) (Hy2 × WF9)	9C, 10
1989	(Hy2 × WF9) (M14 × Oh29)	9C, 10
1990	(Hy2 × WF9) (M14 × Oh43)	9C, 10
1991	(Cl03 × B10) (WF9 × Oh7A)	9C, 10
1992	(Cl03 × B14) (WF9 × Oh7A)	9C, 10
1993	(WF9 × Oh41) (B10 × B14)	9C, 10
1994	(Cl03 × WF9) (Oh29 × Oh41)	9C, 10
1995	(Hy2 × Oh7) (38-11 × Oh41)	9C, 10
1996	(Cl03 × B14) (Hy2 × Oh7)	9C, 10
1997	(Cl03 × Oh41) (Hy2 × Oh7)	9C, 10
1998	(M14 × 187-2) (WF9 × B6)	2C
1999	(Cl03 × Oh43) (M14 × WF9)	2C
3005	(M14 × WF9) (B14 × W64A)	2C
3006	(M14 × W64A) (WF9 × B14)	2C
3007	(R161 × WF9) (R169 × B14)	2C
3008	(R165 × WF9) (R168 × B14)	2C
3009	(B14 × B21) (A297 × W64A)	2C
3010	(Cl03 × N24) (WF9 × B14)	6C
3011	(Cl03 × Oh43) (WF9 × B14)	6C
3012	(Cl03 × B37) (Oh28 × Oh43)	6C
3013	(Cl03 × Oh41) (Hy2 × WF9)	6C
3014	(Hy2 × WF9) (B14 × Oh41)	6C
3015A	(WF9 × B14) (B37 × N24)	6C
3016	(WF9 × B14) (B37 × Oh43)	2C, 6C
3017	(WF9 × B14) (B37 × Oh45)	6C
3018	(WF9 × B14) (B38 × N24)	6C
3019 (Iowa 4880)	(WF9 × B14) (B38 × Oh43)	6C
3020	(WF9 × B14) (N6 × Oh43)	6C

(Table is continued on next page)

Table 22.—Continued

Hybrid	Pedigree	Table No.
Illinois hybrids (continued)		
3021	(WF9 × B14) (N6 × Oh45)	6C
3022	(WF9 × B14) (N22A × Oh43)	6C
3023A	(WF9 × B14) (N24 × Oh43)	6C
3024	(WF9 × B14) (N24 × Oh422)	6C
3025	(WF9 × B14) (N610 × Oh43)	6C
3026	(WF9 × B14) (N610 × Oh45)	6C
3027	(WF9 × B14) (N611 × Oh43)	6C
3029	(WF9 × B14) (Oh43 × Oh45)	6C
3030	(WF9 × B14) (Oh43 × Oh422)	6C
3032	(WF9 × B38) (Oh28 × Oh43)	6C
3033	(WF9 × B40) (Oh28 × Oh43)	6C
3034	(WF9 × N6) (Oh28 × Oh43)	6C
3035	(WF9 × N613) (Oh28 × Oh43)	6C
3036	(B14 × N6) (Oh28 × Oh43)	6C
3037	(B14 × B40) (Oh28 × Oh43)	6C
3038	(B37 × Oh26A) (Oh28 × Oh43)	6C
3039	(B37 × B38) (Oh28 × Oh43)	6C
3040	(B35 × B40) (Oh28 × Oh43)	6C
3041	(WF9 × B35) (Oh28 × Oh43)	6C
3042	(WF9 × B14) (B40 × Oh45)	6C
3043	(R71 × R109B) (WF9 × B14)	2C, 6C
3044	(R109B × R113) (WF9 × B14)	2C, 6C
3045	(R109B × R168) (WF9 × B14)	2C, 6C
3046	(R113 × R168) (WF9 × B14)	2C, 6C
3047	(R71 × R113) (WF9 × B14)	2C, 6C
3048	(R71 × R168) (WF9 × B14)	2C, 6C
3049	(Hy2 × WF9) (R71 × R109B)	9C
3050	(Hy2 × WF9) (R109B × R113)	9C
3051	(Hy2 × WF9) (R109B × R168)	9C
3052	(Hy2 × WF9) (R113 × R168)	9C
3054	(R109B × R113) (WF9 × 38-11)	9C
3055	(R109B × R168) (WF9 × 38-11)	9C
3056	(R113 × R168) (WF9 × 38-11)	9C
3057	(R71 × R109B) (R113 × R168)	2C, 3D, 9C, 14B, 17C
3058	(R71 × R113) (R109B × R168)	3D, 9C, 17C
3059	(R71 × R168) (R109B × R113)	3D, 9C, 14B, 17C
3060	(R129 × R159) (R166 × R168)	9C
3061	(R129 × R159) (R168 × R169)	9C
3062	(R159 × R161) (R168 × R169)	9C
3063	(R159 × R163) (R165 × R168)	9C
3064	(R159 × R163) (R166 × R168)	9C
3065	(R159 × R163) (R168 × R169)	9C
3066	(R159 × R168) (R163 × R165)	9C
3067	(R159 × R169) (R161 × R168)	9C
3069	(R71 × R101) (R105 × R129)	9C
3070	(R71 × R105) (R163 × R168)	9C
3071	(R71 × R129) (R101 × R105)	9C
3072	(R71 × R129) (R105 × R168)	9C
3073	(R71 × R163) (R105 × R168)	9C
3074	(R71 × R168) (R105 × R163)	9C
3075	(Hy2 × WF9) (R95 × R101)	9C
3076	(Hy2 × WF9) (R96 × R101)	9C
3077	(Hy2 × WF9) (R96 × B36)	9C
3078	(Hy2 × WF9) (R96 × Oh451)	9C
3079	(Hy2 × WF9) (R101 × 38-11)	9C

(Table is continued on next page)

Table 22.—Continued

Hybrid	Pedigree	Table No.
Illinois hybrids (continued)		
3080	(Hy2 × WF9) (R101 × Oh451)	9C
3081	(Hy2 × WF9) (R109B × R127)	9C
3082	(Hy2 × WF9) (R109B × B38)	9C
3083	(Hy2 × WF9) (R109B × K720)	9C
3084	(Hy2 × WF9) (R127 × B38)	9C
3085	(Hy2 × WF9) (R127 × L317)	9C
3086	(Hy2 × WF9) (R127 × K720)	9C
3087	(Hy2 × WF9) (R127 × K721)	9C
3088	(Hy2 × WF9) (R127 × N25)	9C
3089	(Hy2 × WF9) (38-11 × Oh451)	9C
3090	(Hy2 × WF9) (B36 × Oh451)	9C
3091	(Hy2 × WF9) (B38 × L317)	9C
3092	(Hy2 × WF9) (B38 × K720)	9C
3093	(Hy2 × WF9) (B38 × N25)	9C
3094	(Hy2 × WF9) (B38 × N35)	9C
3095	(Hy2 × WF9) (L317 × K720)	9C
3096	(R74 × R101) (R129 × WF9)	9C
3097	(R95 × R101) (WF9 × 38-11)	9C
3098	(R98 × R101) (WF9 × 38-11)	9C
3100	(R101 × N12) (WF9 × 38-11)	9C
3101	(R101 × N23) (WF9 × 38-11)	9C
3102	(R101 × Oh41) (WF9 × 38-11)	9C
3103	(R109B × R154) (WF9 × 38-11)	9C
3104	(R109B × N25) (WF9 × 38-11)	9C
3105	(R129 × R154) (WF9 × 38-11)	9C
3106	(R129 × N25) (WF9 × 38-11)	9C
3107	(R154 × B38) (WF9 × 38-11)	9C
3108	(R154 × K721) (WF9 × 38-11)	9C
3109	(R154 × K722) (WF9 × 38-11)	9C
3110	(R154 × N25) (WF9 × 38-11)	9C
3111	(R159 × R163) (R168 × WF9)	9C
3112	(WF9 × 38-11) (B38 × N25)	9C
3113	(WF9 × 38-11) (K722 × N25)	9C
3114	(Hy2 × WF9) (R101 × Cl.38B)	9C
3115	(R127 × N35) (WF9 × 38-11)	9C
3116	(R127 × K721) (WF9 × 38-11)	9C
3117	(R127 × R154) (WF9 × 38-11)	9C
3118	(Hy2 × WF9) (38-11 × B38)	9C
3119	(Hy2 × WF9) (R154 × B38)	9C
3120	(Hy2 × WF9) (R127 × 38-11)	9C
3121	(Hy2 × WF9) (R127 × R154)	9C
3124	(Hy2 × WF9) (R71 × R168)	6C, 9C
3125	(R71 × R168) (WF9 × 38-11)	9C
3126	(R101 × Mo3) (38-11 × K201)	17C
3127	(38-11 × K201) (Mo3 × Mo8)	17C
3128	(38-11 × K201) (Mo3 × Mo9)	17C
3129	(R101 × Mo8) (38-11 × K201)	17C
3130	(R101 × Mo9) (38-11 × K201)	17C
3131	(R129 × Mo3) (38-11 × K201)	17C
3132	(38-11 × K201) (Mo8 × Mo9)	17C
3133	(R127 × Mo3) (38-11 × K201)	17C
3135	(R71A × Mo3) (38-11 × K201)	17C
3136	(R74 × R101) (38-11 × K201)	17C
3137	(38-11 × K201) (Mo4 × Mo9)	17C

(Table is continued on next page)

Table 22.—Continued

Hybrid	Pedigree	Table No.
Illinois hybrids (concluded)		
3138	(R129 × Mo9) (38-11 × K201)	17C
3139	(R71A × R101) (38-11 × K201)	17C
3140	(38-11 × K201) (Ky126 × Cl.21E)	17C
3141	(38-11 × K201) (K763 × Ky126)	17C
3142	(38-11 × K201) (Ky126 × Ok11)	17C
3143	(38-11 × K201) (Ky126 × Oh7B)	17C
3144	(38-11 × K201) (K711 × Ok11)	17C
3145	(R129 × Mo9150) (38-11 × K201)	17C
3146	(R118 × Mo9150) (38-11 × K201)	17C
3147	(R118 × R129) (38-11 × K201)	17C
3148	(R74 × Mo9150) (38-11 × K201)	17C
3149	(R74 × R129) (38-11 × K201)	17C
3150	(R74 × R118) (38-11 × K201)	17C
3151	(WF9 × 38-11) (B14 × Oh41)	9C
3152	(M14 × WF9) (B14 × Oh43)	2C
3159	(M14 × 187-2) (WF9 × Oh43)	3D
3160	(WF9 × Oh7) (B14 × Oh43)	6C
3169C	(WF9 × B37) (Oh28 × Oh43)	6C
2235W	(H21 × K64) (33-16 × Mo2RF)	17C
2246W	(R144 × R145) (R148 × R149)	9C, 10, 17ABC
2247W	(R144 × R145) (R146 × R148)	2ABC, 6ABC
6016	(R78 × K4) (R84 × 38-11)	11A
6021	(R75 × R76) (R84 × K4)	11A
6052	(R78 × 38-11) (R84 × K4)	11A
6062	(R76 × K4) (R78 × R84)	11A
6075	(R75 × R83) (R78 × R87)	11A
6076	(R76 × R78) (R87 × R117)	11A
6084	(R78 × R117) (R84 × R87)	11A
Miscellaneous hybrids		
AES 510	(WF9 × W22) (H19 × B9)	2ABC
AES 610	(M14 × A73) (Oh43 × Oh51A)	2ABC, 3D, 4B, 14B
AES 702 (Ill. 1790)	(C103 × M14) (Hy2 × WF9)	2ABC, 6ABC, 7B, 9BC, 14B
AES 805 (Ill. 1770)	(C103 × Oh45) (WF9 × 38-11)	6ABC, 9ABC, 13B, 14B, 17ABC, 18B
AES 806	(Hy × WF9) (N6 × N15)	6ABC, 9ABC
AES 808	(WF9 × 38-11) (H14 × Oh43)	9ABC
AES 903W	(H28 × K55) (H30 × K41)	17ABC
AES 904W	(K64 × Mo22) (T111 × T115)	17ABC
A-101	(Hy2 × Oh7) (88-4A × SS101)	9C
A-102	(Hy2 × Oh7) (128-4A × SS101)	9C
GCP 6220	(WF9 × 38-11) (O-1265 × GT107)	9C, 17C
Ind. 4655	(C103 × Oh43) (P8 × WF9)	9BC
Ind. 4656	(P8 × WF9) (H14 × Oh43)	9BC
Ind. 5409	(M14 × B14) (WF9 × W22)	2BC
Ind. 5655	(WF9 × 38-11R) (Oh7B × Oh45)	9BC
Ind. 6225 (Ill. 1959)	(M14 × W64A) (B14 × A297)	2C
Ind. 6615	(H49 × H55) (H53 × B14)	17C
Ind. 6623	(C103 × H53) (WF9 × H52)	9C
Ind. 6833	(WF9 × H52) (H54 × H60)	9C
Ind. 6874	(H49 × H52) (H59 × H60)	17C
Iowa 4297	(M14 × 187-2) (WF9 × I.205)	6ABC
Iowa 4757	(M14 × WF9) (B16 × Oh51A)	2BC
Iowa 4779 (Ill. 1862)	(M14 × WF9) (Oh43 × Oh51A)	2BC
Iowa 4809	(M14 × WF9) (B14 × B37)	6BC
Iowa 4879 (Ill. 3016A)	(WF9 × Oh43) (B14 × B37)	6BC

(Table is concluded on next page)

Table 22. — Concluded

Hybrid	Pedigree	Table No.
Miscellaneous hybrids (concluded)		
Iowa 4880 (Ill. 3019)	(WF9 × Oh43) (B14 × B38)	6C
Iowa 4906	(WF9 × B7) (B14 × B39)	9C
Iowa 4907	(WF9 × B7) (B10 × B14)	9C
Iowa 4912 (Ill. 3014A)	(Hy × Oh41) (WF9 × B14)	9BC
ISP 2	(CI03 × Oh45) (M14 × WF9)	2ABC
Kan. 1884	(Hy × R59) (N6 × Oh28)	9C
Kan. 2472W	(H30 × K41) (K55 × K697)	17C
Mich. 52-25	(M14 × WF9) (MS212 × Oh51A)	2BC
Mich. 53-151	(WF9 × MS209) (MS106 × MS107)	2BC
Mich. 54-70	(WF9 × MS107) (MS208 × Oh43)	2C
Mich. 54-116	(WF9 × MS120) (MS211 × Oh43)	2C
Minn. CB4603 (Ill. 3002)	(B14 × A297) (A295 × W64A)	2BC
Minn. CB4621	(B14 × A239) (A295 × W64A)	2BC
Mo. 800-3	(K201 × T202) (Mo9284 × Cl.21E)	17C
Mo. 804 (Ill. 1445)	(38-11 × Cl.21E) (K4 × Cl.7)	17ABC, 18B
Mo. 916	(Mo9108 × Cl.21E) (Oh7B × Oh29)	17BC
Mo. 958	(B41 × Oh7A) (Mo3 × Cl.21E)	17C
Mo. 4060AW	(H30 × K41) (Mo9187W × N72)	9BC
Nebr. 1924	(Hy × WF9) (B14 × N6)	6BC
Ohio M15	(A × W23) (Oh26 × Oh51)	2C
Ohio K24	(WF9 × Oh51A) (Oh33 × Oh40B)	2ABC
Ohio 5317 (CB4726A) (Ill. 3028)	(WF9 × B14) (Oh28 × Oh43)	6BC
TRF 3W	(4413-2 × 4417-1) (K55 × K64)	17C
TRF 9W	(4413-2 × 4417-1) (K6 × K55)	17C
TRF 10	(4401-19 × 4408-6) (4409-6 × Oh29)	17C
TRF 13W	(4413-2 × 4908-2) (K55 × 4914-2)	17C
U.S. 13	(Hy × L317) (WF9 × 38-11)	9ABC, 11B, 12D, 17ABC
U.S. 523W	(K55 × K64) (Ky27 × Ky49)	17ABC
U.S. 619W	(K55 × Cl.64) (Ky27 × Ky49)	17C





UNIVERSITY OF ILLINOIS-URBANA

Q 630 .7L68 C001
BULLETIN. URBANA

621-640 1957-59



3 0112 019528790